



ECOLOGICAL NICHE MODELING AND AN UPDATED GEOGRAPHICAL DISTRIBUTION MAP OF *Leptodactylus notoaktites* HEYER, 1978 (ANURA, LEPTODACTYLIDAE) WITH A NEW OCCURRENCE RECORD

Guilherme de Toledo Figueiredo^{1*}, Diego José Santana²,
Luis Fernando Storti³ & Luiz dos Anjos⁴

¹ Universidade Estadual de Londrina, Departamento de Biologia Animal e Vegetal, Programa de Pós-Graduação em Ciências Biológicas, Rodovia Celso Garcia Cid, PR 445 Km 380, CEP 86057-970, Londrina, PR, Brazil.

² Universidade Federal do Mato Grosso do Sul, Centro de Ciências Biológicas e da Saúde, Av. Costa e Silva s/nº, CEP 79070-900, Campo Grande, MS, Brazil.

³ Samauma Consultoria Ambiental Ltda, Rua Jericó, n. 159, apt. 163, CEP 05435-040, São Paulo, SP, Brazil.

⁴ Universidade Estadual de Londrina, Departamento de Biologia Animal e Vegetal, Rodovia Celso Garcia Cid, PR 445 Km 380, CEP 86057-970, Londrina, PR, Brazil.

E-mails: figueiredoguidetol@gmail.com (*corresponding author); santanadiegojose@yahoo.com; anurofaunalfs@gmail.com; llanjos@sercomtel.com.br

Abstract: *Leptodactylus notoaktites* Heyer, 1978 (Anura, Leptodactylidae) is a Neotropical frog that can be found in open areas, forest edges, and inside forest clearings in southern Brazil. In this study, we present an updated distribution map of this species for a variety of vegetation types in the Atlantic Rain Forest and report a new occurrence record in the Alto Paraná Atlantic Forests. We also performed an Ecological Niche Modeling (ENM) which combined environmental variables with occurrence records to predict environmentally suitable areas for this species. Our study was based on published data, specimens collected in the field, and specimens from Brazilian herpetological collections. The ENM predicted high environmental suitability ranging from São Paulo, Paraná, and Rio Grande do Sul states, mainly in Serra do Mar Coastal Forest areas, while the lowest values were in the states of Rio de Janeiro, Minas Gerais, Mato Grosso do Sul and some inland regions in the states of São Paulo and Paraná. Based on the distribution of *L. notoaktites*, we suggest that field efforts should be extended to inland regions of the Atlantic Rain Forest. In fact, species restricted to coastal regions of the Atlantic Rain Forest could have larger ranges than expected if data from such inland regions was available.

Keywords: advertisement call; anurans; Atlantic Rain Forest; environmental variables; Paraná state.

INTRODUCTION

Leptodactylus notoaktites Heyer, 1978 (Anura, Leptodactylidae) belongs to the *L. mystaceus* species complex, which is placed in the *L. fuscus* species group (de Sá *et al.* 2014). This species is a

medium-sized frog (mean snout-vent length = 47.4 mm) characterized by an upper shank and tibial barred pattern, two distinct dorsolateral folds, posterior thigh with light stripe, some individuals have white tubercles on the sole of the foot (Figure 1), and only individuals with a light mid-dorsal

stripe have six dorsolateral folds (Heyer 1978). The species presents a white lip and black rostral range from the snout to the eardrum (Forlani *et al.* 2010). As for most species in the *L. fuscus* group, *L. notoaktites* lays its eggs in foam nests built in underground burrows, and as the pond floods the exotrophic tadpoles develop (Haddad & Prado 2005).




Figure 1. Vouchered adult male alive (a) of *Leptodactylus notoaktites* (Anura, Leptodactylidae) from the Parque Estadual Mata São Francisco, between Cornélio Procópio and Santa Mariana municipalities, state of Paraná, Brazil. Photos of the preserved specimen as follows: (b) dorsal and (c) ventral views; (d) detail of the sole of foot showing presence of white tubercles (white arrow) and (e) detail of posterior thigh with light stripe (white arrows). Photo (a) by Luis Fernando Storti; (b to e) by Guilherme de Toledo Figueiredo.

Leptodactylus notoaktites is found in open habitats along rivers, forest edges, and clearings inside forests. Such clearings inside forests are natural or the result of anthropogenic disturbance (Skuk & Heyer 2004). Currently, the species is distributed in the states of São Paulo, Paraná, and Santa Catarina, mainly along the coast up to 900 m

a.s.l., which includes several protected areas (Skuk & Heyer 2004, de Sá *et al.* 2014). Despite being classified by the Red List of Endangered Species of International Union for Conservation of Nature as Least Concern and the fact that the species has a stable population trend, potential threats to this species include habitat loss due to deforestation, advance of agricultural areas, and infrastructure development for tourism, which makes additional research on the species geographical distribution necessary (Skuk & Heyer 2004).

Ecological Niche Modeling (ENMs) are important tools when evaluating species range, as they provide robust predictions of distributions or suitable environmental regions for species. Given their strong dependence on environmental variables (Duellman & Trueb 1994), ENMs seem particularly valid for anurans. ENMs use environmental variable associations and known species occurrences regions to generate models that define abiotic conditions where survival and reproduction are possible for the populations (Guisan & Thuiller 2005). Furthermore, these models can be used to define cryptic species (Raxworthy *et al.* 2007), predict species with potential invasion success (Peterson 2003, Peterson *et al.* 2006), maintain rare or endangered species (Engler *et al.* 2004), determine impacts of climate changes (Wiens *et al.* 2009), determine priority conservation areas (Chen 2009), and model the spread of crop pests (Venette *et al.* 2010).

Currently, there are several methods to generate ENMs from a set of environmental variables such as BIOCLIM (Nix 1986), MARS (Friedman 1991), DOMAIN (Carpenter *et al.* 1993), GARP (Elith *et al.* 2006), and Maxent (Phillips *et al.* 2006). However, in some particular situations uncertainties in model predictions can arise (Araújo & New 2007). Most of such limitations are due to the fact that ENMs use only presence and/or absence data, which can create a narrower true distribution than expected (Sinclair *et al.* 2010, Vasconcelos *et al.* 2012). If ENMs could include information on biotic interactions or restrictions then dispersal models would become more realistic. To avoid this limitation, it is possible to establish a priori criteria to build the ENMs. Priori criteria include: (I) potentially or abiotically suitable occurrence areas of an organism in a

future climate change scenario, and (II) areas with suitable environmental conditions for organisms with little geographical distribution information (Vasconcelos & Nascimento 2016, Vasconcelos *et al.* 2017).

Herein, we provide an updated distribution map and new occurrence record of *L. notoaktites* with comments on its distribution across several types of vegetation in the Atlantic Rain Forest, based on published data, a specimen collected by the authors, and specimens housed in Brazilian herpetological collections. With these information, we generated a map that indicates areas with potentially suitable environmental conditions for *L. notoaktites* using an ENM approach.

MATERIAL AND METHODS

Species data collection

A single calling male of *L. notoaktites* (MZUEL-1575; snout-vent length = 49.85 mm) was collected on October 23, 2011 at the Parque Estadual Mata São Francisco (PEMSF), a forest remnant of 865 ha located between the municipalities of Cornélio Procópio and Santa Mariana, state of Paraná, Brazil. To confirm its identity, we recorded the specimen advertisement call (air temperature 26°C) at night in an underground burrow at the forest's edge. Calls were recorded using a SONY ICD PX-820 digital recorder coupled with a Yoga HT-81 microphone and analyzed with a sampling frequency of 44 kHz and sample size of 16 bits. We analyzed the calls with Raven Pro 1.5 for Macintosh (Cornell 2014) and constructed audiospectrograms in R software using the package "seewave" (Sueur *et al.* 2008; R Development Core Team 2014) with the following parameters: FFT window width = 256, Frame = 100, Overlap = 75, and flat top filter. Terminology used in the call description follows Köhler *et al.* (2017). For morphological and acoustic comparisons with all calls from the *L. mystaceus* species complex (Table 1), we used data from Heyer *et al.* (1996) and de Sá *et al.* (2014). Four advertisement calls of the collected male in this study were deposited at FNJV (Fonoteca Neotropical "Jacques Vielliard" of Universidade Estadual de Campinas) sound collection (FNJV-33018 to 33021). Another individual was analyzed at the Museu de Zoologia

of Universidade Estadual de Londrina (MZUEL-1333), collected by M. Z. de Lima on April 15, 2009 at Parque Ecológico Klabin, in the municipality of Telêmaco Borba and mentioned in the unpublished thesis of Machado (2004).

All data based on previously published studies (see Table 1) were obtained by searching in the database of "Scientific Electronic Library Online" (Scielo; www.scielo.org) and "Google Scholar" (www.scholar.google.com) on March 2016 with the following search terms: "*Leptodactylus notoaktites*" and "*Leptodactylus notoaktites* distribution". We collected data from the SpeciesLink Database (Cria 2016) and analyzed specimens housed in the following Brazilian herpetological collections: CFBH (Coleção Célio Fernando Baptista Haddad, Universidade Estadual Paulista, Rio Claro, São Paulo); DZSJRP-Amphibia-adults (Coleção de Anfíbios DZSJRP, Universidade Estadual Paulista, São José do Rio Preto, São Paulo); MCP-Anfibios (Coleção de Anfíbios, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Rio Grande do Sul); SinBiota (Sistema de Informação do programa Biota /FAPESP, Fundação de Amparo à Pesquisa do Estado de São Paulo, Campinas, São Paulo), ZUEC-AMP(Coleção de Anfíbios do Museu de Zoologia da UNICAMP, Campinas, São Paulo), UFMG-AMP (Coleção de Anfíbios do Centro de Coleções Taxonômicas da Universidade Federal de Minas Gerais, Belo Horizonte, Minas Gerais); FNJV (Fonoteca Neotropical "Jacques Vielliard", Universidade Estadual de Campinas, Campinas, São Paulo); and MZUEL (Coleção de Herpetofauna do Museu de Zoologia da Universidade Estadual de Londrina, Londrina, Paraná). We only considered records those identified with the term "*Leptodactylus notoaktites*". Records containing taxonomic inaccuracies (e.g., cf., aff. and gr.) were not considered. Vegetation types in the distribution map of *L. notoaktites* followed the ecoregions of Olson *et al.* (2001).

Environmental data collection

We used variables associated with the biological conditions necessary for the occurrence of the taxon: (i) 19 bioclimatic variables available at WorldClim (Fick & Hijmans 2017), which are already well documented in the literature as important bioclimatic variables for amphibian

distributions (e.g., Duellman & Trueb 1994); (ii) altitude, slope and aspect. Altitude is highly correlated with temperature and humidity (Hoersch *et al.* 2002) and with solar radiation that plays an important role on moisture for habitat selection by amphibians (Wyman 1988). Slope and aspect were related to a proxy for the amount of solar radiation on the ground surface (Blank & Blaustein 2012); (iii) vegetation type (biome characteristics), which is related to species habitat (Toledo & Batista 2012); and (iv) percent of tree cover (MODIS), which affects amphibian species composition and distribution (Figueiredo *et al.*, unpublished data; Skelly *et al.* 1999). Bioclimatic variables were obtained from WorldClim (WorldClim database version 2.0, <http://www.worldclim.org/>) and were interpolated to 30 arc-sec resolution (Fick & Hijmans 2017) with WGS84 projection; altitude, slope, aspect, and vegetation type from National Aeronautics and Space Administration - NASA (<http://www2.jpl.nasa.gov/srtm/>; Amaral *et al.* 2013); and percent of tree cover from Global Landcover Facility (<http://glcf.umd.edu/data/>; Amaral *et al.* 2013).

Model building and evaluations

We used the 35 records (see Table 1) to generate the ENM based on the machine-learning modeling method Maxent (maximum entropy algorithm; Phillips *et al.* 2006), which estimates areas with potentially suitable environmental conditions for *L. notoaktites*. In this approach the niche suitability ranges from zero to one; the closer to one, the greater environmental suitability for the species. To avoid over-prediction and low specificity for species distribution (*i.e.*, Amazonian Rainforest), we cropped the bioclimatic layers to span from latitude -18 to -35 and longitude -38 to -58 (values in decimal degrees). To avoid model over-parameterization, we removed multicollinearity variables ($r > 0.8$) and variables with low contribution to the model (< 1%) using a Jackknife test of variable importance determined by their biological relevance for *L. notoaktites*. We used this approach because the test excluded one variable at a time when running the model (Baldwin 2009) and, therefore, provided information about the activity of each variable in the model, showing how important each variable

was in explaining the species distribution and how much unique information each variable contained. This approach can also present highly correlated variables, thereby allowing us to determine if the rate of contribution values are constrained due to these multicollinear variables (Phillips 2017). Nine of the 19 original bioclimatic variables, vegetation type, and percent of tree cover were used in the final model: Bio2 (Mean Diurnal Range); Bio4 (Temperature Seasonality); Bio5 (Max temperature warmest month); Bio9 (Mean temperature driest quarter); Bio14 (Precipitation of Driest Month); Bio15 (Precipitation Seasonality); Bio17 (Precipitation driest quarter); Bio18 (Precipitation of Warmest Quarter) and Bio19 (Precipitation coldest quarter). The evaluation and performance of species distribution model was tested using the threshold-independent receiver operation characteristic (ROC) analysis. The area under the ROC curve (AUC) ranged from 0 to 1. After tenfold cross validation we used the model test of the AUC scores (Elith *et al.* 2006, Philips *et al.* 2006). AUC values greater than 0.9 are considered very good; AUC between 0.7 and 0.9 are good; and AUC values less than 0.7 are uninformative (Swets 1988). All the analyses were performed in the R platform vs. 3.3.2 (R Core Team 2014), MaxEnt software v. 3.4.1 (Phillips *et al.* 2006) and implemented in ArcGIS 10.5 for desktop (Esri 2016).

RESULTS

A total of 35 records were compiled: field collection (1); SpeciesLink database records (17) and literature records (17). Based on these records we found the distribution of *L. notoaktites* in three states of Brazil (Table 1 and Figure 2): São Paulo (16 records), followed by Paraná (12), and Santa Catarina (7). Most of the records are found in the coastal range of these states. Regarding the vegetation type where the records are located we listed six types (Figure 2): Alto Paraná Atlantic Forests (7); Araucaria Moist Forest (7); Grasslands (5); Serra do Mar Coastal Forests (14); and Southern Atlantic Mangroves (2).

The advertisement call of the collected specimen of *L. notoaktites* was composed of a single, harmonic, unpulsed note, with duration of




Figure 2. Geographical distribution map of *Leptodactylus notoaktites* (Anura, Leptodactylidae) in the southern and southeastern Brazil with indication of vegetation types. States: SP = São Paulo, PR = Paraná, SC = Santa Catarina. Records: White dots = literature records, white squares = SpeciesLink records, and white triangles = new records. Localities: 1. Parque Estadual Mata São Francisco; 2. Telêmaco Borba; 3. Benedito Novo; 4. Blumenau; 5. Brusque; 6. Jaraguá do Sul; 7. Corupá; 8. São Bento do Sul; 9. Piraquara; 10. Antonina; 11. Parque Estadual Ilha do Cardoso; 12. Iguape; 13. Ribeirão Branco; 14. Eldorado; 15. Registro; 16. Campinas; 17. Rio Claro; 18. Corumbataí; 19. Rio negro; 20. Serra Dona Francisca; 21. Fazenda Rio Grande; 22. Guaratuba; 23. Morretes; 24. São José dos Pinhais; 25. Reserva Natural Salto Morato; 26. Barra do turvo; 27. Iporanga; 28. PETAR; 29. Parque Estadual Intervales; 30. Parque Estadual Carlos Botelho; 31. Ribeirão Grande; 32. Capão Bonito; 33. Campos Gerais; 34. Piraí do Sul; 35. Curitiba.

0.074 – 0.091 s (0.080 ± 0.005 s; Table 2 and Figure 3). The call had ascending frequency modulation, with a dominant frequency of 689 – 1205 Hz (1029 ± 214 Hz; Table 2 and Figure 3). Hence, the advertisement call analyzed confirms the identity of the specimen collected in the PEMSF as *L. notoaktites*. AUC test for the replicate runs of the ENMs generated for *L. notoaktites* (Figure 4) was 0.99 ± 0.004 , indicating a very good model. The environmental suitability was explained primarily by Temperature Seasonality (bio4; 34.3%), Precipitation of Warmest Quarter (bio18; 23.8%) and Precipitation of Driest Month (bio14; 13.8%).

Altitude, slope, and aspect were discarded due to their poor contribution to the model (< 1%). The highest environmental suitability in the model (> 0.5; represented by the water blue and blue colors in Figure 4) ranged from São Paulo, Paraná, Santa Catarina, and Rio Grande do Sul states, whereas the most probable areas of occurrence were along the coastal range of São Paulo, Paraná, and Santa Catarina states. The lowest values (< 0.3; green color in Figure 4) were in the states of Rio de Janeiro, Minas Gerais, Mato Grosso do Sul, and some inland regions of São Paulo and Paraná states.


Table 1. Known localities for *Leptodactylus notoaktites* (Anura, Leptodactylidae) in southern and southeastern Brazil. Id = Identification number included in Figure 2. States: SP= São Paulo; PR = Paraná; SC= Santa Catarina. Source type: F = Field; LT = Literature; SL = Records from SpeciesLink database.

Id	Municipalities (Locality)	State	Latitude	Longitude	Source Type	Reference
1	Cornélio Procópio; Santa Mariana (Parque Estadual Mata São Francisco)	PR	23.1590°S	50.5660°W	F	Present study (MZUEL 1575)
2	Telêmaco Borba (Fazenda Monte Alegre)	PR	24.2857°S	50.4170°W	LT	Machado (2004); (MZUEL 1333)
3	Benedito Novo (Reserva Biológica Sassafrás)	SC	26.7826°S	49.3640°W	SL	Present study (UFMG-AMP 3091)
4	Blumenau (Parque das Nascentes)	SC	26.9193°S	49.0660°W	SL	Present study (PUCRS-MCP 8681)
5	Brusque (RPPN Chácara Edith)	SC	27.0982°S	48.9170°W	SL	Present study (PUCRS-MCP 7639)
6	Jaraguá do Sul (Vila Nova - Vila Lenzi)	SC	26.4860°S	49.0660°W	SL	Present study (PUCRS-MCP 1482)
7	Corupá (RPPN Emílio Fiorentino Battistella)	SC	26.4250°S	49.2430°W	SL	Present study (CFBH 28718)
8	São Bento do Sul (Rio Vermelho - Natal)	SC	26.2500°S	49.3780°W	SL	Present study (UFMG-AMP 9866)
9	Piraquara (Barragem Piraquara)	PR	25.4410°S	49.0630°W	SL	Present study (CFBH 11037)
10	Antonina (Reserva Natural Rio da Cachoeira)	PR	25.4280°S	48.7110°W	SL	Present study (CFBH 12361)
11	Cananéia (Parque Estadual Ilha do Cardoso - Charco do Haras)	SP	25.1380°S	47.9670°W	SL	Present study (FNJV 12960)
12	Iguape (Estação Ecológica da Juréia - Itatins)	SP	24.7000°S	47.5500°W	SL	Present study (ZUEC-AMP 16904)
13	Ribeirão Branco (Fazenda do João Zaqueu e Mathedi)	SP	24.2200°S	48.7650°W	SL	Present study (CFBH 6896)
14	Eldorado (Pousada Recanto das Águas)	SP	24.5200°S	48.1000°W	SL	Present study (CFBH 10665)
15	Registro (Vale do Ribeira)	SP	24.4875°S	47.8432°W	SL	Present study (SINBIOTA C10166T86634)
16	Campinas (Barão Geraldo, estrada da Rhodia)	SP	22.9000°S	47.0603°W	SL	Present study (ZUEC-AMP 20973)
17	Rio Claro (Horto Florestal)	SP	22.4100°S	47.5600°W	SL	Present study (CFBH 4335)
18	Corumbataí (Sítio Santa Amélia)	SP	22.2200°S	47.6254°W	SL	Present study (DZSJRP 5663)
19	Rio Negro (Parque Municipal São Luiz de Tolosa)	PR	26.0847°S	49.8060°W	LT	Santos & Conte (2014)
20	Joinville (Serra Dona Francisca)	SC	26.1950°S	49.0350°W	LT	Mariotto (2014)
21	Fazenda Rio Grande (Fazenda Gralha Azul)	PR	25.9280°S	49.1980°W	LT	Conte & Rossa-Feres (2007)
22	Guaratuba (Colônia Castelhanos)	PR	25.8820°S	48.5740°W	LT	Cunha <i>et al.</i> (2010)
23	Morretes (Condomínio Rio Sagrado)	PR	25.5650°S	48.7990°W	LT	Armstrong & Conte (2010)
24	São José dos Pinhais (Serro e Gemido)	PR	25.4105°S	49.0300°W	LT	Conte & Rossa-Feres (2006)

Table 1. Continued on next page...

Table 1. ...continued

25	Guaraqueçaba (Reserva Natural Salto Morato)	SP	25.1686°S	48.2984°W	LT	Garey & Hartmann (2012)
26	Barra do Turvo (Parque Estadual Jacupiranga)	SP	24.7561°S	48.5042°W	LT	Aguiar-de-Domenico (2008)
27	Iporanga (Parque Estadual da Caverna do Diabo)	SP	24.5330°S	48.7000°W	LT	Aguiar-de-Domenico (2008)
28	Apiaí; Iporanga (Parque Estadual Turístico do Alto Ribeira PETAR)	SP	24.4952°S	48.6471°W	LT	Araújo <i>et al.</i> (2010)
29	Ribeirão Grande (Parque Estadual Intervales)	SP	24.3232°S	48.2800°W	LT	Bertoluci & Rodrigues (2002)
30	Tapiraí; São Miguel Arcanjo; Capão Bonito (Parque Estadual Carlos Botelho)	SP	24.1414°S	47.9740°W	LT	Forlani <i>et al.</i> (2010)
31	Ribeirão Grande (Fazenda Intermontes)	SP	24.0900°S	48.3600°W	LT	Tacioli (2012)
32	Capão Bonito (Fazenda Intervales)	SP	24.0000°S	48.3400°W	SL	Present study (ZUEC-AMP 8525)
33	Ponta Grossa; Carambeí; Castro (Campos Gerais)	PR	25.0500°S	49.9500°W	LT	Crivellari <i>et al.</i> (2014)
34	Piraí do Sul (Flona de Piraí do Sul)	PR	24.5660°S	49.9160°W	LT	Foerster (2014)
35	Curitiba (Parques Municipais de Curitiba)	PR	25.4160°S	49.2500°W	LT	Crivellari <i>et al.</i> (2014)

**Figure 3.** Audiospectrogram and oscilogram of the advertisement call of *Leptodactylus notoaktites* (Anura, Leptodactylidae) recorded from a single male at Parque Estadual Mata São Francisco, Paraná state, Brazil. (air temperature 26°C; snout-vent length = 49.85 mm).

DISCUSSION

Regarding the geographic distribution of *L. notoaktites* (see Figure 2), our record in the PEMSF, Paraná state, considerably expanded the distribution of the species to the north of the state, and represents a new record for the Alto Paraná Atlantic Forests, an Atlantic Rain Forest vegetation type. The Atlantic Rain Forest was one of the largest rainforests of the Americas with high environmental heterogeneity (Ribeiro *et al.* 2009), being considered a hotspot for conservation (Myers *et al.* 2000). The PEMSF is located approximately 258 km in a straight-line from the type locality of the species in the municipality of Iporanga, São Paulo state, which consists of Serra do Mar Coastal Forests, and is about 140 km in a straight-line from the nearest record in the Telêmaco Borba municipality, located in the Araucaria Moist Forest (Machado 2004).

Other species similar to *L. notoaktites*, which include species from the *L. mystaceus* complex (*L. cupreus*, *L. didymus*, *L. elenae*, *L. mystaceus*, and *L. spixii*), have distinct geographical distributions. *Leptodactylus mystaceus*, which occurs in

Amazonia, and Minas Gerais and São Paulo states in mesic enclaves of northeastern Brazil, as well as northern portions of the Atlantic Rain Forests of Brazil (de Sá *et al.* 2014), is the only species of the *L. mystaceus* complex that occurs in the nearby region of our new record of *L. notoaktites* in the PEMSF (see Affonso *et al.* 2011). Nevertheless, when comparing the morphology of both species it is clear that only individuals of *L. notoaktites* with a light mid-dorsal stripe also have a pair of dorsal folds, while *L. mystaceus* lacks both a pair of dorsal folds and a light mid-dorsal stripe (de Sá *et al.* 2014). Besides, there are several acoustic differences between the species. For instance, the advertisement call of *L. mystaceus* presents no harmonic structure, while the advertisement call of *L. notoaktites* is harmonically structured (de Sá *et al.* 2014, present study). Therefore, bioacoustic studies can be useful for understanding ecological processes under climate change (Møller 2010) or anthropic actions in fragile systems (deforestation and fragmentation of habitats) (Tucker *et al.* 2014), fostering new perspectives in ecology and conservation fields described as Ecoacoustics (Sueur & Farina 2015).

Table 2. Acoustic parameter comparisons of the advertisement call for the *Leptodactylus mystaceus* (Anura, Leptodactylidae) species complex. *Species with frequency modulations in the beginning/end of call.

Species	Dominant Frequency (kHz)	Call Duration (s)	Pulses	Call rate	Harmonic	Reference
<i>L. notoaktites</i>	689–1.205 Hz/1.550–1.722 Hz*	0.074–0.091s	No	0.6/s	Yes	Present study
<i>L. cupreus</i>	2800–3058 Hz	0.16s	No	–	Yes	Caramaschi <i>et al.</i> (2008)
<i>L. didymus</i>	510–1.510 Hz	0.09–0.32s	Yes	1.4–3.1/s	Yes	Heyer <i>et al.</i> (1996), Köhler & Lötters (2002)
<i>L. elenae</i>	700–870 Hz/1.370–1.500 Hz*	–	No	64–120/min	Yes/No	Barrio (1965), Heyer & Heyer (2002)
<i>L. mystaceus</i>	700–1.400 Hz	0.2s	Yes	1.8/s	No	Heyer (1978), Heyer <i>et al.</i> (1996)
<i>L. spixii</i>	1.500–1.722/1.981–2.067 Hz*	0.12 ± 10s	No	80–97/min	Yes	Bilate <i>et al.</i> (2006)

Most of the records of *L. notoaktites* are along the coastal region of Brazil, mainly in the Serra do Mar Coastal Forests (e.g., Reserva Natural Salto Morato, municipality of Guaraqueçaba, Paraná state; Parque Estadual Intervales, municipalities of Ribeirão Grande, Guapiara, Sete Barras, Eldorado,

and Iporanga, São Paulo state) and Araucaria Moist Forest (e.g., São José dos Pinhais, municipality of Piraí do Sul, Paraná state) (Table 1 and Figure 2). The other records are in the Alto Paraná Atlantic Forest (e.g., Parque Estadual Mata São Francisco, municipality of Rio Claro, Paraná

state) and Grasslands (in the phytogeographic unit of Campos Gerais, municipalities of Ponta Grossa, Carambeí, Castro, Paraná state), where there are ecotone with Grasslands and Araucaria Moist Forest (Table 1 and Figure 2). Some individuals were recorded in Southern Atlantic Mangroves (Estação Ecológica da Juréia – Itatins, municipality of Iguape; Parque Estadual Ilha do Cardoso,

municipality of Cananéia; and Reserva Natural Rio da Cachoeira, municipality of Antonina, São Paulo state). This species also occurs in ecotones with Alto Paraná Atlantic Forests and Cerrado domain vegetation (*e.g.*, municipality of Corumbataí, São Paulo state) and perhaps may even occur in the Cerrado, since the model predicted suitable environments there (see Figure 4).




Figure 4. Predicted distribution of *Leptodactylus notoaktites* (Anura, Leptodactylidae) based on environmental suitability. Occurrence localities can be found in Table 1 and Figure 2.

Records of *L. notoaktites* outside of its occurrence region predicted by ENM may be the result of the lack of field efforts in some regions. This gap in records could limit more robust predictions by the model. This may occur because the geographical range of any species is based on several complex interactions between intrinsic factors of the species (*e.g.*, life history, dispersal abilities, environmental requirements) and extrinsic factors (*e.g.*, variations in space and time), which limit the known distributions (Brown *et al.* 1996). However, many other factors could be

affecting their distribution, such as species colonizing regions outside of their known occurrence and breeding in areas that are significantly different from their former pristine breeding habitats as result of deforestation (Rubbo & Kiesecker 2005). Another possibility is that the species has historically occurred in these environments (in another age) and may now be isolated in refugia (Reside *et al.* 2014). Refugia are habitats where populations of species can retreat to, persist in, and even expand their geographical range over ecological and evolutionary time scales

of millennia (Keppel *et al.* 2012).

Herein we show that the distribution of *L. notoaktites* is not restricted to the Brazilian Coast and that it can also occur in inland forests of the Atlantic Rain Forest. Nevertheless, this species still does not have a large distribution (*e.g.*, *L. mystaceus*), which reinforces the need to maintain suitable habitats for this species for conservation and management actions. For biological conservation, ENMs have been mainly applied to discover biodiversity, study species invasion, conservation efforts, and climate changes effects (Rangel & Loyola 2012). Thus, studies that model species distributions along ecosystems, especially for those species with restricted distribution, can serve as a tool for decision-making and indicate priority sites to be preserved. These regions must have suitable conditions for the survival/persistence of rare and/or endangered species or even species with restricted distribution. Furthermore, we highlight that the gaps in field efforts in certain regions, as in the northern Paraná state, should be considered to estimate the range of *L. notoaktites*. As found for this species, it is possible that several anuran species with known ranges restricted to the Serra do Mar Coastal Forests and Southern Atlantic Mangroves of Brazil could actually have larger distributions. In fact, the anurans diversity in Alto Paraná Atlantic Forests and Araucaria Moist Forests may be higher than expected.

ACKNOWLEDGEMENTS

We are grateful to Instituto Ambiental do Paraná (IAP) and Parque Estadual Mata São Francisco (PEMSF) for providing all infrastructure. ICMBio/SISBIO for collection permits 15363-1. GTF, LFS and DJS thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the scholarship and GTF thanks China Three Georges Corporation for doctoral scholarship.

REFERENCES

Affonso, I. D. P., Delariva, R. L., & Navarro, M. P. 2011. Amphibia, Anura, Leptodactylidae,

- Leptodactylus mystaceus* (Spix, 1824): distribution extension. Check List, 7(2), 198–199. DOI: 10.15560/7.2.198
- Amaral, S., Costa, C. B., Arasato, L. S., Ximenes, A. C., & Rennó, C. D. 2013. AMBDATA: variáveis ambientais para Modelos de Distribuição de Espécies (MDEs). In: M. L. Ribeiro, T. G. Santos, & S. J. Sant'Anna (Eds.), Anais XV Simpósio Brasileiro de Sensoriamento Remoto. pp. 6930–6937. São José dos Campos, SP: Instituto Nacional de Pesquisas Espaciais.
- Aguiar-de-Domenico, E. 2008. Herpetofauna do mosaico de unidades de conservação do Jacupiranga (SP). Master thesis. Departamento de Zoologia da Universidade de São Paulo. p. 200. DOI: 10.11606/D.41.2008.tde-25092008-170838
- Araújo, C. D. O., Condez, T. H., Bovo, R. P., Luiz, A. M., Florestal, H., Vista, B., & Claro, R. 2010. Amphibians and reptiles of the Parque Estadual Turístico do Alto Ribeira (PETAR), SP: an Atlantic Forest remnant of Southeastern Brazil. Biota Neotropica, 10(4), 257–274. DOI: 10.1590/S1676-06032010000400031
- Araújo, M. B., & New, M. 2007. Ensemble forecasting of species distributions. Trends in Ecology & Evolution, 22(1), 42–47. DOI: 10.1016/j.tree.2006.09.010
- Armstrong, C. G., & Conte, C. E. 2010. Taxocenose de anuros (Amphibia: Anura) em uma área de Floresta Ombrófila Densa no Sul do Brasil. Biota Neotropica, 10(1), 39–46.
- Barrio, A. 1965. Afinidades del canto nupcial de las especies cavícolas del género *Leptodactylus* (Anura, Leptodactylidae). Physis, 25(70), 401–410.
- Baldwin, R. A. 2009. Use of maximum entropy modeling in wildlife research. Entropy, 11(4), 854–866. DOI: 10.3390/e11040854
- Bertoluci, J., & Rodrigues, M. T. 2002. Utilização de hábitats reprodutivos e micro-habitats de vocalização em uma taxocenose de anuros (Amphibia) da Mata Atlântica do sudeste do Brasil. Papéis Avulsos de Zoologia 42(11), 287–297. DOI: 10.1590/S0031-10492002001100001
- Bilate, M., Wogel, H., Weber, L. N., & Abrunhosa, P. A. 2006. Vocalizações e girino de *Leptodactylus spixii* Heyer, 1983 (Amphibia, Anura, Leptodactylidae). Arquivos do Museu Nacional, Rio de Janeiro, 64(3), 235–245.

- Blank, L., & Blaustein, L. 2012. Using ecological niche modeling to predict the distributions of two endangered amphibian species in aquatic breeding sites. *Hydrobiologia*, 693(1), 157–167. DOI: 10.1007/s10750-012-1101-5
- Brown, J., Stevens, G., & Kaufman, D. 1996. The Geographic range: size, shape, boundaries, and internal structure. *Annual Review of Ecology and Systematics*, 27(1), 597–623. DOI: 10.1146/annurev.ecolsys.27.1.597
- Caramaschi, U., Feio R. N., & São-Pedro, V. A. 2008. A new species of *Leptodactylus fitzingeri* (Anura, Leptodactylidae) from Serra do Brigadeiro, State of Minas Gerais, southeastern Brazil. *Zootaxa*, 1861, 44–54. DOI: 10.5281/zenodo.183691
- Carpenter, G., Gillison, A. N., & Winter, J. 1993. DOMAIN: a flexible modelling procedure for mapping potential distributions of plants and animals. *Biodiversity & Conservation*, 2(6), 667–680. DOI: 10.1007/BF00051966
- Chen, Y. H. 2009. Conservation biogeography of the snake family Colubridae of China. *North-Western Journal of Zoology*, 5(2), 251–262.
- Conte, C. E., & Rossa-Feres, D. C. 2006. Diversidade e ocorrência temporal da anurofauna (Amphibia, Anura) em São José dos Pinhais, Paraná, Brasil. *Revista Brasileira de Zoologia*, 23(1), 162–175. DOI: 10.1590/S0101-81752006000100008
- Conte, C. E., & Rossa-Feres, D. C. 2007. Riqueza e distribuição espaço-temporal de anuros em um remanescente de Floresta de Araucária no sudeste do Paraná. *Revista Brasileira de Zoologia*, 24(4), 1025–1037. DOI: 10.1590/S0101-81752007000400020
- Cornell Bioacoustics Research Program. 2014. Raven Pro: interactive sound analysis software (version 1.5). The Cornell Lab of Ornithology, Ithaca, New York, USA. Retrieved from <http://www.birds.cornell.edu/brp/raven/Rave nOverview.html>
- Cria - Centro de Referência em Informação Ambiental. 2016. SpeciesLink - Eletronic Database. Retrieved September 05, 2016, from <http://splink.cria.org.br/>
- Crivellari, L. B., Leivas, P. T., Leite, J. C. M., Gonçalves, D. S., Mello, C. M., Rossa-Feres, D. C., & Conte, C. E. 2014. Amphibians of grasslands in the state of Paraná, southern Brazil (Campos Sulinos). *Herpetology Notes*, 7, 639–654.
- Cunha, A. K., Oliveira, I. S., & Hartmann, M. T. 2010. Anurofauna da Colônia Castelhanos, na área de proteção ambiental de Guaratuba, serra do mar Paranaense, Brasil. *Biotemas*, 23(2), 123–134. DOI: 10.5007/2175-7925.2010v23n2p123
- de Sá, R. O., Grant, T., Camargo, A., Heyer, W. R., Ponssa, M. L., & Stanley, E. 2014. Systematics of the Neotropical genus *Leptodactylus* Fitzinger, 1826 (Anura: Leptodactylidae): phylogeny, the relevance of non-molecular evidence, and species accounts. *South American Journal of Herpetology*, 9(s1), s1–s100. DOI: 10.2994/SAJH-D-13-00022.1
- Duellman, W. E., & Trueb, L. 1994. Biology of amphibians. Baltimore and London: The Johns Hopkins University Press: p. 670.
- Elith, J., Graham, C. H., Anderson, R. P., Dudík, M., Ferrier, S., Guisan, A., Hijmans, R. J., Huettmann, F., Leathwick, J. R., Lehmann, A., Li, J., Lohmann, L. G., Loiselle, B. A., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J. McC. M., Townsend Peterson, A., Phillips, S. J., Richardson, K., Scachetti-Pereira, R., Schapire, R. E., Soberón, J., Williams, S., Wisz, M. S., & Zimmermann, N. E. 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, 29(2), 129–151. DOI: 10.1111/j.2006.0906-7590.04596.x
- Engler, R., Guisan, A., & Rechsteiner, L. 2004. An improved approach for predicting the distribution of rare and endangered species from occurrence and pseudo absence data. *Journal of Applied Ecology*, 41(2), 263–274. DOI: 10.1111/j.0021-8901.2004.00881.x
- Esri. 2016. ArcGIS (version 10.5). Environmental Systems Research Inst., Redlands, CA.
- Fick, S. E., & Hijmans, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, 7(12), 4302–4315. DOI: 10.1002/joc.5086
- Foerster, N. E. 2014. Partilha acústica, uso do sítio de vocalização e influência da heterogeneidade ambiental em uma taxocenose de anuros em um remanescente de floresta ombrófila mista. Master thesis. Departamento de Zoologia da

- Universidade Federal do Paraná. p. 70
- Forlani, M. C., Bernardo, P. H., Haddad, C. B. F., & Zaher, H. 2010. Herpetofauna of the Carlos Botelho State Park, São Paulo State, Brazil. *Biota Neotropica*, 10(3), 265–308. DOI: 10.1590/S1676-06032010000300028
- Friedman, J. H. 1991. Multivariate adaptive regression splines. *The Annals of Statistics*, 19(1), 1–67.
- Garey, M. V., & Hartmann, M. T. 2012. Anurans of Reserva Natural Salto Morato, municipality of Guaraqueçaba, State of Paraná, southern Brazil. *Biota Neotropica*, 12(4), 137–145. DOI: 10.1590/S1676-06032012000400015
- Guisan, A., & Thuiller, W. 2005. Predicting species distribution: offering more than simple habitat model. *Ecology Letters*, 8, 993–1009. DOI: 10.1111/j.1461-0248.2005.00792.x
- Haddad, C. F. B., & Prado, C. P. A. 2005. Reproductive modes in frogs and their unexpected diversity in the Atlantic forest of Brazil. *BioScience*, 55(3), 207–217. DOI: 10.1641/0006-3568(2005)055[0207:RMIFAT]2.0.CO;2
- Heyer, W. R. 1978. Systematics of the *fuscus* group of the frog genus *Leptodactylus* (Amphibia, Leptodactylidae). *Natural History Museum of Los Angeles City, Science Bulletin*, 29, 1–85.
- Heyer, R. W., García-Lopez, J. M., & Cardoso, A. J. 1996. Advertisement call variation in the *Leptodactylus mystaceus* species complex (Amphibia: Leptodactylidae) with a description of a new sibling species. *Amphibia-Reptilia*, 17(1), 7–31. DOI: 10.1163/156853896X00252
- Heyer, W. R., & Heyer, M. M. 2002. *Leptodactylus elenae* catalogue of American amphibians and reptiles. *Society for the Study of Amphibians and Reptiles*, 742, 1–5.
- Hoersch, B., Braun, G., & Schmidt, U. 2002. Relation between landform and vegetation in alpine regions of Wallis, Switzerland. A multiscale remote sensing and GIS approach. *Computers, Environment and Urban Systems*, 26(2), 113–139. DOI: 10.1016/S0198-9715(01)00039-4
- Keppel, G., Van Niel, K. P., Wardell-Johnson, G. W., Yates, C. J., Byrne, M., Mucina, L., Schut, A. G. T., Hooper, S. D., & Franklin, S. E. 2012. Refugia: identifying and understanding safe havens for biodiversity under climate change. *Global Ecology and Biogeography*, 21(4), 393–404. DOI: 10.1111/j.1466-8238.2011.00686.x
- Köhler, J., & Lötters, S. 2002. *Leptodactylus didymus*. In: R. Márquez., I. J. de la Riva, J. Bosch, & E. Matheu (Eds.), *Guía sonora de las ranas y sapos de Bolivia*. pp. 32 and CD2, track 11. Madrid: Museo Nacional de Ciencias Naturales, Fonoteca Zoológica.
- Köhler, J., Jansen, M., Rodríguez, A., Kok, P. J. R., Toledo, L. F., Emmrich, M., Glaw, F., Haddad, C. F. B., Rodel, M. O., & Vences, M. 2017. The use of bioacoustics in anuran taxonomy: theory, terminology, methods and recommendations for best practice. *Zootaxa*, 4251(1), 1–124. DOI: 10.11646/zootaxa.4251.1.1
- Machado, R. A. 2004. Ecologia de assembleias de anfíbios anuros no município de Telêmaco Borba, Paraná, sul do Brasil. Doctoral thesis. Departamento de Zoologia da Universidade Federal do Paraná. p. 113.
- Mariotto, L. R. 2014. Anfíbios de um gradiente altitudinal em Mata Atlântica. Master thesis. Departamento de Zoologia da Universidade Federal do Paraná. p. 174.
- Møller, A. P. 2010. When climate change affects where birds sing. *Behavioral Ecology*, 22(1), 212–217. DOI: 10.1093/beheco/arp200
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A., & Kent, J. 2000. Biodiversity hotspots for conservation priorities. *Nature*, 403(6772), 853. DOI: 10.1038/35002501
- Nix, H. 1986. A biogeographic analysis of Australian Elapid snakes. In: R. Longmore (Ed.), *Snakes: atlas of Elapid snakes of Australia*. pp. 4–15. Canberra: Bureau of Flora and Fauna.
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wetengel, W. W., Hedao, P., & Kassem, K. R. 2001. Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience*, 51(11), 933–938. DOI: 10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2
- Peterson, A. T. 2003. Predicting the geography of species' invasions via ecological niche modeling. *The Quarterly Review of Biology*, 78(4), 419–433. DOI: 10.1086/378926
- Peterson, A. T., Papes, M., Reynolds, M. G., Perry,

- N. D., Hanson, B., Regnery, R. L., Hutson, C. L., Muizniek, B., Damon, I. K., & Carroll, D. S. 2006. Native-range ecology and invasive potential of *Cricetomys* in North America. *Journal of Mammalogy*, 87(3), 427–432. DOI: 10.1644/05-MAMM-A-133R3.1
- Phillips, S. J., Anderson, R. P., & Schapire, R. E. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190(3), 231–259. DOI: 10.1016/j.ecolmodel.2005.03.026
- Phillips, S. J. 2017. A brief tutorial on Maxent. Retrieved June 05, 2017, from: http://biodiversityinformatics.amnh.org/open_source/maxent/
- Rangel, T. F., & Loyola, R. D. 2012. Labeling ecological niche models. *Natureza & Conservação*, 10(2), 119–126. DOI: 10.4322/nat.con.2012.030
- Raxworthy, C. J., Ingram, C. M., Rabibisoa, N., & Pearson, R. G. 2007. Applications of ecological niche modeling for species delimitation: a review and empirical evaluation using day geckos (*Phelsuma*) from Madagascar. *Systematic Biology*, 56(6), 907–923. DOI: 10.1080/10635150701775111
- R Core Team. 2014. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Retrieved from <http://www.R-project.org/>
- Reside, A. E., Welbergen, J. A., Phillips, B. L., Wardell-Johnson, G. W., Keppel, G., Ferrier, S., Williams, S. E., & VanDerWal, J. 2014. Characteristics of climate change refugia for Australian biodiversity. *Austral Ecology*, 39(8), 887–897. DOI: 10.1111/aec.12146
- Ribeiro, M. C., Metzger, J. P., Martensen, A. C., Ponzoni, F. J., & Hirota, M. M. 2009. The Brazilian Atlantic Forest: how much is left, and how is the remaining forest distributed? Implications for conservation. *Biological Conservation*, 142(6), 1141–1153. DOI: 10.1016/j.biocon.2009.02.021
- Rubbo, M. J., & Kiesecker, J. M. 2005. Amphibian breeding distribution in an urbanized landscape. *Conservation Biology*, 19(2), 504–511. DOI: 10.1111/j.1523-1739.2005.000101.x
- Santos, E. J., & Conte, C. E. 2014. Riqueza e distribuição temporal de anuros (Amphibia: Anura) em um fragmento de Floresta Ombrófila Mista. *Iheringia, Série Zoologia*, 104(3), 323–333.
- Sinclair, S., White, M., & Newell, G. 2010. How useful are species distribution models for managing biodiversity under future climates? *Ecology and Society*, 15(1), 8.
- Skelly, D. K., Werner, E. E., & Cortwright, S. A. 1999. Long-term distributional dynamics of a Michigan amphibian assemblage. *Ecology*, 80(7), 2326–2337. DOI: 10.1890/0012-9658(1999)080[2326:LTDDOA]2.0.CO;2
- Skuk, G., & Heyer, R. 2004. *Leptodactylus notoaktites*. The IUCN Red List of Threatened Species 2004. Retrieved on March 23, 2016, from <http://dx.doi.org/10.2305/IUCN.UK.2004.RLTS.T57150A11592295.en>
- Sueur, J., Aubin, T., & Simonis, C. 2008. Seewave, a free modular tool for sound analysis and synthesis. *Bioacoustics*, 18(2), 213–226. DOI: 10.1080/09524622.2008.9753600
- Sueur, J., & Farina, A. 2015. Ecoacoustics: the ecological investigation and interpretation of environmental sound. *Biosemiotics*, 8(3), 493–502. DOI: 10.1007/s12304-015-9248-x
- Swets, J. A. 1988. Measuring the accuracy of diagnostic systems. *Science*, 240(4857), 1285–1293. Retrieved on April 15, 2017, from <http://www.jstor.org/stable/1701052>
- Tacioli, A. 2012. Padrão temporal de atividade de vocalização da anurofauna de dois ambientes aquáticos em uma área de compensação ambiental, Serra de Paranapiacaba, (SP). Master thesis. Universidade Estadual Paulista. p. 91.
- Toledo, L. F., & Batista, R. F. 2012. Integrative study of Brazilian anurans: geographic distribution, size, environment, taxonomy, and conservation. *Biotropica*, 44(6), 785–792. DOI: 10.1111/j.1744-7429.2012.00866.x
- Tucker, D., Gage, S. H., Williamson, I., & Fuller, S. 2014. Linking ecological condition and the soundscape in fragmented Australian forests. *Landscape Ecology*, 29(4), 745–758. DOI: 10.1007/s10980-014-0015-1
- Vasconcelos, T. S., Rodriguez, M. A., & Hawkins, B. A. 2012. Species distribution modelling as a macroecological tool: a case study using New World amphibians. *Ecography*, 35(6), 539–548. DOI: 10.1111/j.1600-0587.2011.07050.x

- Vasconcelos, T. S., & do Nascimento, B. T. 2016. Potential climate-driven impacts on the distribution of generalist treefrogs in South America. *Herpetologica*, 72(1), 23–31. DOI: 10.1655/HERPETOLOGICA-D-14-00064
- Vasconcelos, T. S., Antonelli, C. P., & Napoli, M. F. 2017. Mutualism influences species distribution predictions for a bromeliad-breeding anuran under climate change. *Austral Ecology*, 42(7), 869–877. DOI: 10.1111/aec.12509
- Venette, R. C., Kriticos, D. J., Magarey, R. D., Koch, F. H., Baker, R. H. A., Worner, S. P., Raboteaux, N. N. G., McKenney, D. W., Dobesberger, E. J., Yemshanov, D., de Barro, P. J., Hutchison, W. D., Fowler, G., Kalaris, T. M., & Pedlar, J. 2010. Pest risk maps for invasive alien species: a roadmap for improvement. *BioScience*, 60(5), 349–362. DOI: 10.1525/bio.2010.60.5.5
- Wiens, J. A., Stralberg, D., Jongsomjit, D., Howell, C. A., & Snyder, M. A. 2009. Niches, models, and climate change: assessing the assumptions and uncertainties. *Proceedings of the National Academy of Sciences*, 106(2), 19729–19736. DOI: 10.1073/pnas.0901639106
- Wyman, R. 1988. Soil acidity and moisture and the distribution of amphibians in five forests of Southcentral New York. *Copeia*, 1988(2), 394–399. DOI: 10.2307/1445879

Submitted: 04 July 2017

Accepted: 21 November 2017

Associate Editor: Ana C. Delciellos