

Record of Sturnira tildae for Pernambuco, Brazil

FIRST RECORD OF *Sturnira tildae* DE LA TORRE, 1859 (CHIROPTERA, PHYLLOSTOMIDAE) FOR THE STATE OF PERNAMBUCO, BRAZIL

Tereza Cristina dos Santos Leal Martins¹, Edson Silva Barbosa Leal², Carlos Henrique Campos Bezerra Neves^{2,3}, Alan Felipe Ferreira³, Lucas Gonçalves Silva¹, Ana Cristina Lauer Garcia^{2*} & Martín Alejandro Montes^{1,3}

¹Universidade Federal Rural de Pernambuco, Departamento de Biologia, Rua Dom Manoel de Medeiros, s/n, Dois Irmãos, CEP 52171-900, Recife, PE, Brazil.

²Universidade Federal de Pernambuco, Centro Acadêmico de Vitória, Rua Alto do Reservatório, s/n, CEP 55608-680, Vitória de Santo Antão, PE, Brazil.

³Universidade Federal da Paraíba, Programa de Pós-Graduação em Ciências Biológicas, Campus Universitário, s/n, CEP 58051-900, João Pessoa, PB, Brazil.

E-mails: dossantos.t.c@gmail.com, edson.leal76@gmail.com, carloscampos.bn@hotmail.com, lifealan@gmail.com, lucas_gonc@yahoo.com.br, alauergarcia@yahoo.com.br (*corresponding author), martin.montes@ufrpe.br

Abstract: Here we present a new record of *Sturnira tildae* in Pernambuco state, Brazil, based on a capture of a pregnant female in an anthropized environment. This finding expands the known distribution for the species in the northern Atlantic Forest. The species nearest record is about 500 km far from our collection locality. Dental and morphometric characteristics were used to identify the species. These features were compared with those of *S. lilium* with whom the species has morphological similarities. *Sturnira tildae* was identified based on its spatulate internal upper incisor teeth, and the lower first and second molars with low lingual cusps and shallow notches. The forearm size of the captured specimen (greater than 45 mm) was also used for identification of *S. tildae*.

Keywords: Atlantic Forest; bat; distribution; morphology; range extension.

Approximately a quarter of the mammal species of the world are bats (Simmons 2005). In the tropical portion of the Neotropics, the diversity of bats can reach 50 % of the mammalian fauna richness (Timm 1994). Brazil is the second country with the highest number of bat species, with 182 species, distributed in nine families and 69 genera (Nogueira *et al.* 2018). Many recent studies have updated the list of Brazilian bat species (*e.g.* Feijó *et al.* 2015, Gregorin *et al.* 2016, Velazco *et al.* 2017, Pavan *et al.* 2018) and reported an increase in geographic distribution for several species in the country (Rocha *et al.* 2017, Lima *et al.* 2018, Nunes *et al.* 2018). Bats of the genus *Sturnira* (Chiroptera, Phyllostomidae) are endemic to the New World and comprise 19 species (Velazco & Patterson 2013, 2019), four of which occur in Brazil: *S. lilium* (Geoffroy 1810), *S. giannae* Velazco & Patterson, 2019, *S. magna* De La Torre, 1966, and *S. tildae* De La Torre, 1959 (Nogueira *et al.* 2014). *Sturnira tildae* is an important disperser of seeds and occurs in the eastern portion of Colombia, southern Venezuela, Guianas, Peru, Bolivia, Trinidad and Brazil (Simmons 2005). In Brazil, the species has been recorded in the Amazon, Atlantic Forest, Cerrado and Caatinga biomes (Paglia *et al.* 2012).

Externally, *S. tildae* does not have facial stripes, has a mask of dark hairs around the eyes, the uropatagium and calcar are much reduced, and the tail is absent. The pelage coloration is usually orange, ranging from yellowish to brownish (Reis *et al.* 2013). In areas where it occurs in sympatry, it differs from *S. lilium* by dental and morphometric characteristics (Simmons & Voss 1988).

Here we present a new record of *S. tildae* in northeastern Brazil, which expands its geographic distribution. Bats were collected in 10 nights of sampling, between August and November 2016, at the Technology Institute of Pernambuco (8°05'27.7"S, 34°55'41.1"W), located in Recife, Pernambuco state. It is inside an urban area, with a mixture of ombrophilous dense Atlantic Forest and exotic vegetation. The climate in the area is warm and humid with an annual

temperature average of 26 °C, with a higher incidence of rainfall between March and August (INMET 2019).

Five mist nets (12 m x 2.5 m) were used in each sampling night. The nets were set from 17h00-00h00 and checked for bats at 20-min intervals. Collections occurred on nights of a new moon and no rain. The captures were authorized by the SISBIO (license nº 20383-1) and by the Animal Ethics Committee of the Federal Rural University of Pernambuco (license nº 067/2015). External and cranial measurements (in mm) were taken using a digital caliper with a capacity of 20 cm and precision of 0.05 mm. The specimen of *S. tildae* was fixed in 10 % formalin and preserved in 70 % ethanol (Papavero 1994). The skull was removed and cleaned according to Pacheco (2004). The following external and cranial measurements were recorded according to Nogueira *et al.* (2008) and Velazco *et al.* (2010): body length, forearm length, ear length, skull length, condylobasal length, postorbital constriction, breadth across upper molar, braincase breadth, zygomatic breadth and maxillary toothrow length. The body mass was verified with a high precision balance (EK5055, max 5 kg, precision of 0.1 g). The specimen was deposited in the Collection of Mammals of the Federal University of Pernambuco, under the number UFPE 3642.

Identification of the *S. tildae* was based on qualitative and quantitative characters according to De La Torre (1959), Marinkelle & Cadena (1971), Simmons & Voss (1998), and Reis *et al.* (2013). Measurements of the external and cranial characters of this specimen were compared with those of *S. lilium* from Brazil (Taddei 1975), Trinidad (Goodwin & Greenhall 1964), French Guiana (Simmons & Voss 1998), and Mexico (Goodwin & MacDougall 1969). In order to update the knowledge about the geographical distribution of *S. tildae*, bibliographic surveys were carried out in books, articles, monographs, dissertations, and theses. The keywords used were *Sturnira*, *Sturnira tildae*, and *S. tildae*.

A total of 117 bats were captured in the present study (Table 1). A single female specimen of *S. tildae*, pregnant (by palpation), non-nursing, was collected on October 09, 2016. The

captured specimen had brown pelage coloration, dark hairs around the eyes, hairs from the ventral region lighter than the dorsal, very short calcar, and no tail (Figure 1). The skull is long, with a small braincase and broad, high rostrum. The specimen was identified as *S. tildae* based on its spatulate internal upper incisor teeth, and the lower first and second molars with low lingual cusps and shallow notches (Marinkelle & Cadena 1971, Simmons & Voss 1998, Reis *et al.* 2013, Figure 2).

Table 1. Species and abundance of bats captured at the Technology Institute of Pernambuco, Recife, Brazil, between August and November 2016.

Taxon	N
Family Phyllostomidae	
Subfamily Carolliinae	
<i>Carollia perspicillata</i> (Linnaeus, 1758)	21
Subfamily Glossophaginae	
<i>Glossophaga soricina</i> (Pallas, 1766)	7
Subfamily Phyllostominae	
<i>Phyllostomus discolor</i> (Wagner, 1843)	4
Subfamily Stenodermatinae	
<i>Artibeus lituratus</i> (Olfers, 1818)	15
<i>A. obscurus</i> (Schinz, 1821)	51
<i>A. planirostris</i> (Spix, 1823)	10
<i>Sturnira lilium</i> (E. Geoffroy, 1810)	5
<i>S. tildae</i> De La Torre, 1959	1
<i>Platyrrhinus lineatus</i> (E. Geoffroy, 1810)	3



Figure 1. *Sturnira tildae* (female, UFPE 3642) captured in Recife, Pernambuco state, Brazil.




Figure 2. Lateral (A) and dorsal perspective (B) of skull, lower jaw (C) and upper incisors (D) of female *Sturnira tildae* (UFPE 3642) captured in Recife, Pernambuco state, Brazil. Bars = 5 mm.

The external and cranial measurements (forearm length, cranial length, condylobasal length, postorbital constriction, zygomatic breadth, and maxillary tooththrow length) as well as the body mass of the captured specimen are according with the description of *S. tildae* by De La Torre (1959), and by Marinkelle & Cadena (1971), Simmons & Voss (1998) and Reis *et al.* (2013) (Table 2). The forearm length is an important taxonomic character used to differentiate *S. tildae* from *S. lilium*. The first species has a forearm length of more than 45 mm while in *S. lilium* this measure does not exceed this value (Taddei 1975, Gannon *et al.* 1989, Simmons & Voss 1998, Reis *et al.* 2013).

Table 2. External and craniodental measurements (in millimeters), and body mass (in grams) for *Sturnira tildae* (female, UFPE 3642) from Recife, Pernambuco state, Brazil. Including mean values observed for *S. tildae* females from Trinidad (De La Torre 1959), French Guiana (Simmons & Voss 1998) and Colombia (Marinkelle & Cadena 1971), and means for *S. lilium* females from Brazil (São Paulo, Taddei 1975), Trinidad (Goodwin & Greenhall 1964), French Guiana (Simmons & Voss 1998) and Mexico (Goodwin & MacDougall 1969).

Measurements	<i>Sturnira tildae</i>				<i>Sturnira lilium</i>			
	Brazil, Pernambuco (UFPE 3642)	Trinidad	French Guiana	Colombia	Brazil, São Paulo	Trinidad	French Guiana	Mexico
External	N = 1	N = 1	N = 13	N = 60	N = 20	N = 3	N = 19	N = 5
Total length	66.5		69.8		61.62		64.5	
Forearm length	45.9	54.1	46.1	47.4	42.42	41.67	41.9	39.76
Ear length	15.5		18.1	15.2	16.10		16.4	
Body mass	26.3		22.5			16.1	19.3	
Craniodental	N = 1	N = 1	N = 5	N = 50	N = 15	N = 3	N = 11	N = 5
Cranial length	23.6	23.3	22.76	23.7	22.11	21.8	21.77	21.34
Condylbasal length	22.6	21.1	21.26	22.3	20.01	20.37	20.05	19.36
Postorbital constriction	6.5	6.3	6.16	6.6	5.95		5.79	5.56
Breadth across upper molars	7.9	8.1	8.21	8.4	8.13		8	
Braincase breadth	10.2		10.8	10	10.41	10.17	10.18	9.76
Zygomatic breadth	14.9	13.9	14.15	14.6	13.86	12.73	13.42	13.12
Maxillary toothrow length	7.6	7.1	6.87	7.1	6.40	6.57	6.46	6.32

This is the fifth record of *S. tildae* in the Northeast region of Brazil and the first in the Pernambuco state. The nearest historical record is about 500 km to the west from Chapada do Araripe, Ceará, in the Caatinga biome (Novaes & Laurindo 2014). The other records in the Northeast region of the country are all from the Atlantic Forest biome of the Bahia state, more than 850 km to the south (Faria 2006, Faria & Baugarten 2007, Table 3 and Figure 3).

Table 3. Location records of *Sturnira tildae* in Brazil. Codes refer to numbers indicated in the Figure 3.

Code	Latitude/ Longitude	Locality/State	Reference
1	1°36'02''N/52°28'59''W	Serra do Navio/Amapá	Martins <i>et al.</i> (2006)
2	1°16'57''N/51°35'03''W	Ferreira Gomes/Amapá	Martins <i>et al.</i> (2006)
3	1°10'05''N/49°33'51''W	Cutias/Amapá	Martins <i>et al.</i> (2011)
4	0°15'55''N/53°06'01''W	Almeirim/Amapá	Martins <i>et al.</i> (2006)
5	0°00'00''N/51°00'00''W	Macapá/Amapá	Martins <i>et al.</i> (2011)
6	2°15'07''S/62°37'55''W	Novo Airão/Amazonas	Barnett <i>et al.</i> (2006)
7	2°24'04''S/59°43'07''W	Preto da Eva/Amazonas	Bernard (2002)
8	2°37'02''S/64°36'54''W	Maraã/Amazonas	Pereira <i>et al.</i> (2009)
9	2°55'09''S/59°52'59''W	Manaus/Amazonas	Capaverde (2015)
10	3°10'10''S/67°23'01''W	Jutaí/Amazonas	Santos (2017)
11	3°05'07''S/59°57'12''W	Manaus/Amazonas	Neves (2017)
12	8°12'17''S/63°52'57''W	Canutama/Amazonas	Silva & Bobrowiec (2015)
13	1°09'01''S/49°37'51''W	Anajás/Pará	Marques-Aguiar <i>et al.</i> (2002)
14	1°13'09''S/48°16'49''W	Santa Bárbara do Pará/Pará	Fonseca (2006)
15	1°16'11''S/49°21'05''W	Muaná/Pará	Marques-Aguiar <i>et al.</i> (2002)
16	2°30'09''S/54°57'17''W	Santarém/Pará	Bernard & Fenton (2002)
17	3°36'11''S/55°44'47''W	Aveiro/Pará	Presley <i>et al.</i> (2008)
18	7°41'18''S/51°52'02''W	Ourilândia do Norte/Pará	Peters <i>et al.</i> (2006)
19	7°27'11''S/73°40'47''W	Mâncio Lima/Acre	Nogueira <i>et al.</i> (2004)

20	10°10'10''S/68°30'09''W	Rio Branco/Acre	Santos <i>et al.</i> (2017)
21	10°03'55''S/67°36'51''W	Senador Guimard/Acre	Silva (2016)
22	7°42'10''S/47°19'06''W	Goiatins/Tocantins	Maas <i>et al.</i> (2018)
23	8°30'11''S/48°27'44''W	Colinas do Tocantins/Tocantins	Maas <i>et al.</i> (2018)
24	10°30'06''S/46°10'16''W	Mateiros/Tocantins	Gregorin <i>et al.</i> (2011)
25	7°21'51''S/39°20'11''W	Barbalha/Ceará	Novaes & Laurindo (2014)
26	8°05'27.7''S/34°55'41.1''W	Recife/Pernambuco	This Study
27	14°47'11''S/39°04'07''W	Ilhéus/Bahia	Faria & Baugarten (2007)
28	15°17'18''S/39°15'11''W	Una/Bahia	Faria (2006)
29	9°04'07''S/60°36'44''W	Colniza/Mato Grosso	Pinto (2008)
30	12°48'48''S/51°46'12''W	Ribeirão Casalheira/Mato Grosso	Pine <i>et al.</i> (1970)
31	13°43'26''S/47°50'13''W	Cavalcante/Goiás	Oliveira (2000)
32	17°51'57''S/51°43'21''W	Jataí/Goiás	Pina <i>et al.</i> (2013)
33	20°26'21''S/54°38'15''W	Campo Grande/Mato Grosso do Sul	Jesus <i>et al.</i> (2015)
34	19°39'08''S/40°07'17''W	Aracruz/Espírito Santo	Peracchi & Albuquerque (1993)
35	22°25'21''S/42°45'47''W	Cachoeiras de Macacu/Rio de Janeiro	Souza <i>et al.</i> (2015)
36	22°22'22''S/44°35'56''W	Itatiaia/Rio de Janeiro	Luz <i>et al.</i> (2013)
37	22°49'28''S/43°31'08''W	Rio de Janeiro/Rio de Janeiro	Menezes <i>et al.</i> (2015)
38	22°59'10''S/44°06'21''W	Mangaratiba/Rio de Janeiro	Luz <i>et al.</i> (2011)
39	23°10'05''S/44°12'03''W	Angra dos Reis/Rio de Janeiro	Esbérard <i>et al.</i> (2006)

40	22°20'28''S/47°50'34''W	Pirassununga/São Paulo	Muylaert <i>et al.</i> (2014)
41	23°20'47''S/46°29'01''W	Mairiporã/São Paulo	Bertola <i>et al.</i> (2005)
42	23°22'22''S/45°50'26''W	Ubatuba/São Paulo	Garbino (2016)
43	23°44'41''S/46°02'07''W	Bertioga/São Paulo	Garbino (2016)
44	24°09'18''S/48°36'21''W	Guapiara/São Paulo	Passos <i>et al.</i> (2003)
45	24°17'55''S/47°00'09''W	Peruíbe/São Paulo	Gimenez & Ferrarezzi (2004)
46	24°33'52''S/48°40'49''W	Iporanga/São Paulo	Garbino (2016)
47	24°13'27''S/48°04'46''W	Sete Barras/São Paulo	Garbino (2016)
48	24°41'13''S/48°00'01''W	Jacupiranga/São Paulo	Garbino (2016)
49	24°42'04''S/47°52'21''W	Pariquera-Açu/São Paulo	Garbino (2016)
50	25°03'15''S/47°52'41''W	Cananéia/São Paulo	Garbino (2016)
51	25°10'06''S/48°18'17''W	Guaraqueçaba/Paraná	Munster (2008)
52	25°27'17''S/48°54'49''W	Morretes/Paraná	Brito (2011)
53	25°25'07''S/48°54'57''W	Morretes/Paraná	Miretzki <i>et al.</i> (2002)
54	27°01'17''S/49°01'06''W	Blumenau/Santa Catarina	Althoff (2007)
55	28°28'43''S/49°15'09''W	Pedras Grandes/Santa Catarina	Carvalho <i>et al.</i> (2013)
56	28°37'03''S/49°01'11''W	Jaguaruna/Santa Catarina	Bôlla <i>et al.</i> (2017)

Sturnira tildae has been mainly observed in preserved environments (Miretzki *et al.* 2006), but the locality where it was captured in our field survey has anthropic alterations, being a small forest fragment inserted in the urban area of Recife. According to Reis *et al.* (2013) there is also a possibility for capturing *S. tildae* in anthropized environments.




Figure 3. Geographic distribution of *Sturnira tildae* in Brazil and the new record (star) in Recife, Pernambuco state. The numbers refer to the codes in Table 3.

In the present study, we observed higher abundance of *S. lillium* compared to *S. tildae* (Table 1), a pattern that is maintained throughout the Atlantic Forest (Faria 2006, Faria & Baumgarten 2007, Brito 2011, Carvalho *et al.* 2013, Luz *et al.* 2013, Pires & Cademartori 2018). In the Atlantic Forest of Northeast Brazil (present study, Faria 2006, Faria & Baumgarten 2007) the abundance of *S. lillium* is lower than in the southern portion of this biome (Passos *et al.* 2003, Esbérard *et al.* 2006, Brito 2011, Carvalho *et al.* 2013, Luz *et al.* 2013, Souza *et al.* 2015). Here we observed *S. tildae* with relative frequency below 1 %, similar to the results recorded throughout the Brazilian Atlantic Forest (Esbérard *et al.* 2006, Faria 2006, Faria & Baumgarten 2007, Luz *et al.* 2011, Carvalho *et al.* 2013; Luz *et al.* 2013, Menezes *et al.* 2015, Souza *et al.* 2015).

The difficulty in identifying *S. tildae* occurs due to its morphological similarities with *S. lilium* (Simmons & Voss 1998). It is very possible that the similarity of the external morphology between these species and the overlapping in their geographic distributions (Sampaio *et al.* 2016, Velazco & Patterson 2017) can lead to identification errors. In this way, it is recommended that mammalogists pay great attention in the identification of *Sturnira* genus, mainly when individuals are captured, identified and released in the field.

REFERENCES

- Althoff, S.L. 2007. A Comunidade de quirópteros, sua biologia e ecologia no Parque Natural Municipal Nascentes do Garcia, Estado de Santa Catarina, Brasil. Doctoral thesis. Universidade Federal do Rio Grande do Sul. p.131.
- Barnett, A.M., Sampaio, E., Kalko, E.K.V., Shapley, R. L., Fischer, E., Camargo, G. & Herrera, B. 2006. Bats of Jaú National Park, central Amazônia, Brazil. *Acta Chiropterologica*, 8(1), 103-128. DOI:10.3161/150811006777070820
- Bernard, E. 2002. Diet, activity and reproduction of bat species (Mammalia, Chiroptera) in Central Amazonia, Brazil. *Revista Brasileira de Zoologia*, 9(1), 173-188. DOI:10.1590/S0101-81752002000100016
- Bernard, E., & Fenton, B. 2011. Species diversity of bats (Mammalia: Chiroptera) in forest fragments, primary forests, and savannas in central Amazonia, Brazil. *Canadian Journal of Zoology*, 80(6), 1124-1140. DOI:10.1139/z02-094
- Bertola, P.B., Aires, C.C., Favorito, S.E., Graciolli, G., Amaku, M., & Pinto-da-Rocha, R. 2005. Bat flies (Diptera: Streblidae, Nycteribiidae) parasitic on bats (Mammalia: Chiroptera) at Parque Estadual da Cantareira, São Paulo, Brazil: parasitism rates and host-parasite associations. *Memórias do Instituto Oswaldo Cruz*, 100(1), 25-32. DOI: 10.1590/S0074-02762005000100005

- Bôlla, D.A.S., Ceron, K., Carvalho, F., Matia, D.L., Luiz, M.R., Panatta, K.A., & Zocche, J.J. 2017. Mastofauna terrestre do sul de Santa Catarina: mamíferos de médio e grande porte e voadores. *Tecnologia e Ambiente*, 23(1), 61-78. DOI: 10.18616/ta.v23i0.3906
- Brito, J.E.C. 2011. Morcegos do Parque Estadual do Pico Marumbi: frugivoria e germinação de sementes. Master thesis. Universidade Federal do Paraná, p.53.
- Capaverde, U.D. 2015. Mudanças sutis na elevação determinam a estrutura das assembleias de morcegos Phyllostomidae na Amazônia central. Master thesis. Instituto Nacional de Pesquisas Amazônicas. p. 51.
- Carvalho, F., Fabián, M.E., & Menegheti, J.O. 2013. Vertical structure of an assemblage of bats (Mammalia: Chiroptera) in a fragment of Atlantic Forest in Southern Brazil. *Zoologia*, 30(5), 491-498. DOI:10.1590/S1984-46702013000500004
- De La Torre, L. 1959. A new species of bat of the genus *Sturnira* (Phyllostomidae) from the island of Trinidad, West Indies. *Chicago Academy of Sciences*, 166(1), 1-6.
- Esbérard, C.E.L., Jordão-Nogueira, T., Luz J. L., Melo, G.G.S., Mangolin, R., Jucá, N., Raíces, D.S. L., Enrici, M.C., & Bergallo, H.G. 2006. Morcegos da Ilha Grande, Angra dos Reis, RJ, Sudeste do Brasil. *Revista Brasileira de Zoociências*, 8(2), 147-153. DOI: 10.34019/2596-3325.2006.v8.24115
- Faria, D. 2006. Phyllostomid bats of a fragmented landscape in the north-eastern Atlantic forest, Brazil. *Journal of Tropical Ecology*, 22, 531-542. DOI: 10.1017/S0266467406003385
- Faria, D., & Baumgarten, J. 2007. Shade cacao plantations (*Theobroma cacao*) and bat conservation in southern Bahia, Brazil. *Biodiversity and Conservation*, 16(2), 291-312. DOI:10.1007/s10531-005-8346-5
- Feijó, A., Rocha, P.A., & Althoff, S.L. 2015. New species of *Histiotus* (Chiroptera: Vespertilionidae) from northeastern Brazil. *Zootaxa*, 4048(3), 412-427. DOI:10.11646/zootaxa.4048.3.4

- Fonseca, R.T.D. 2006. Diversidade da quiropterofauna (Mammalia) no Parque Ecológico de Gunma, Santa Bárbara do Pará. Master thesis. Universidade Federal do Pará. p.107.
- Garbino, G.S.T. 2016. Research on bats (Chiroptera) from the state of São Paulo, southeastern Brazil: annotated species list and bibliographic review. *Arquivos de Zoologia*, 47(3), 43-128. DOI:10.11606/issn.2176-7793.v47i3p43-128
- Gannon, M.R. Willig, M.R., & Jones, J.K. 1989. *Sturnira lilium*. *Mammalian Species*, 333(1), 1-5. DOI: 10.2307/3504237
- Gimenez, E.A., & Ferrarezzi, H. 2004. Diversidade de morcegos no sudeste da Mata Atlântica. In: O.A.V. Marques & W. Duleba (Eds.), *Estação Ecológica Juréia-Itatins: ambiente físico, flora e fauna*. pp. 314-330. Ribeirão Preto: Holos.
- Goodwin, G.G., & Greenhall, A.M. 1964. New records of bats from Trinidad and comments on the status of *Molossus trinitatus* Goodwin. *American Museum Novitates*, 2195, 1-23.
- Goodwin, G.G., & MacDougall, T.B. 1969. Mammals from the State of Oaxaca, Mexico, in the *American Museum of Natural History*. *Bulletin of the American Museum of Natural History*, 141(1), 1-318.
- Gregorin, R., Moras, L.M., Acosta, L.H., Vasconcellos, K.L., Poma, J.L., Santos, F.R., & Paca, R.C. 2016. A new species of *Eumops* (Chiroptera: Molossidae) from southeastern Brazil and Bolivia. *Mammalian Biology*, 81(3), 235-246. DOI: 10.1016/j.mambio.2016.01.002
- Gregorin, R., Gonçalves, E.A., Caroline, C., & Carmignotto, A.P. 2011. Morcegos (Mammalia: Chiroptera) da Estação Ecológica Serra Geral do Tocantins: composição específica e considerações taxonômicas. *Biota Neotropica*, 11(1), 299-311. DOI:10.1590/S1676-06032011000100028
- INMET. 2019. Instituto Nacional de Meteorologia (Retrieved on April 15th, 2019, from <http://www.inmet.gov.br/portal>).
- Jesus, C.C., Torres, J.M., & dos Anjos, E.A. 2015. Plantas consumidas por morcegos na Lagoa da Cruz, Campo Grande, MS. *Multitemas*, 45(1), 123-136.

- Lima, A.C.S., Cardoso, F.H.S., Mendes, S.B., Fraga, E.C., & Barros, M.C. 2018. New records of Niceforo's big-eared bat, *Trinycteris nicefori* (Sanborn, 1949) (Chiroptera, Phyllostomidae), from the state of Maranhão, Brazil. *ZooKeys*, 787(1), 127-134. DOI: 10.3897/zookeys.787.26538
- Luz, J.L., Costa, L.M., Jordão-Nogueira, T., Esbérard, C.E.L., & Bergallo, H. G. 2013. Morcegos em área de Floresta Montana, Visconde de Mauá, Resende, Rio de Janeiro. *Biota Neotropica*, 13(2), 190-195. DOI: 10.1590/S1676-6032013000200018
- Luz, J.L., Costa, L.M., Lourenço, E.C., & Esbérard, C.E.L. 2011. Morcegos (Mammalia, Chiroptera) da Reserva Rio das Pedras, Rio de Janeiro, sudeste do Brasil. *Biota Neotropica*, 11(1), 95-102. DOI:10.1590/S1676-06032011000100009
- Maas, A.C.S., Gomes, L., Martins, M.A., Dias, D., Pol, A., Chaves, G.F., Schutte, M., Araújo, R., & Peracchi, A.L. 2018. Bats in a Cerrado landscape of Northern Brazil: species occurrence, influence of environmental heterogeneity and seasonality, and eight new records for the state of Tocantins. *Mammalia*, 82(5), 469-480. DOI: 10.1515/mammalia-2017-0023
- Marinkelle, C.J., & Cadena, A. 1971. Remarks on *Sturnira tildae* in Colombia. *Journal of Mammalogy*, 52 (1), 235-237. DOI: 10.2307/1378460
- Marques-Aguiar, S.A., Melo, C.C.S., Aguiar, G.F.S., & Queiroz, J.A.L. 2002. Levantamento preliminar da mastofauna da região de Anajás-Muaná, Ilha de Marajó, Pará, Brasil. *Revista Brasileira de Zoologia*, 19(3), 841-854. DOI:10.1590/S0101-81752002000300022.
- Martins, A.C.M., Bernard, E., Gregorin, R. 2006. Inventários biológicos rápidos de morcegos (Mammalia, Chiroptera) em três unidades de conservação do Amapá, Brasil. *Revista Brasileira de Zoologia*, 23(4), 1175-1184. DOI:10.1590/S0101-81752006000400026.
- Martins, A.C.M., Bernard, E., Gregorin, R., Da Silva, W.A.S. 2011. Filling data gaps on the diversity and distribution of Amazonian bats (Chiroptera), the case of Amapá, easternmost Brazil. *Zoologia*, 28(2), 177-185. DOI:10.1590/S1984-46702011000200004.

- Menezes, L.F., Pinto, A.C.D.C., Contildes, M.D.R., & Peracchi, A.L. 2015. Lista de morcegos (Mammalia Chiroptera) do Parque Natural Municipal da Serra do Mendanha, Município do Rio de Janeiro, RJ, Brasil. In: J.A.L. Pontes (Ed.), Biodiversidade carioca. Rio de Janeiro, Technical Books: pp. 238-245.
- Miretzki, M., Perrachi, A.L., & Bianconi, G.V. 2002. Southernmost records of *Sturnira tildae* de La Torre, 1959 (Chiroptera: Phyllostomidae) in Brazil. *Mammalia*, 66(2), 306-308.
- Miretzki, M. 2006. Padrões de distribuição de mamíferos na Floresta Atlântica brasileira. Doctoral thesis. Universidade de São Paulo. p. 294.
- Munster, C.L. 2008. Dieta de morcegos frugívoros (Chiroptera, Phyllostomidae) na Reserva Natural do Salto Morato. Monography. Universidade Federal do Paraná. p. 30.
- Muylaert, R.L., Teixeira, R.C., Hortenci, L., Estêvão, J.R., Rogeri, P.K., Mello, M.A.R. 2014. Bats (Mammalia: Chiroptera) in a cerrado landscape in São Carlos, southeastern Brazil. *Check List*, 10(2), 287-291. DOI:10.15560/10.2.287.
- Neves, L.R. 2017. Efeito de borda sobre a estrutura de comunidade de morcegos em floresta de terra-firme, Amazônia Central. Master thesis. Universidade Federal do Amazonas. p. 36.
- Nogueira, M.R., Fabio, S.P., & Peracchi, A.L. 2004. Gastrointestinal helminth parasitism in fruit-eating bats (Chiroptera, Stenodermatinae) from western Amazonian Brazil. *Revista de Biologia Tropical*, 52(2), 387-92. DOI:10.15517/rbt.v52i2.15254.
- Nogueira, M.R., Lima, I.P., Moratelli, R., Tavares, V.C., Gregorin, R., & Peracchi, A.L. 2014. Checklist of Brazilian bats, with comments on original records. *Check List*, 10(4), 808–821. DOI:10.15560/10.4.808
- Nogueira, M.R., Lima, I.P., Garbino, G.S.T., Moratelli, R., Tavares, V.C., Gregorin, R., & Peracchi A.L. 2018. Updated checklist of Brazilian bats: version 2018.1. Comitê da Lista de Morcegos do Brasil—CLMB. Sociedade Brasileira para o Estudo de Quirópteros (SBEQ). (Retrieved on August 15th, 2019, from www.sbeq.net/updatelist).

- Nogueira, M.R., Pol, A., Monteiro, L.R., & Peracchi, A.L. 2008. First record of Miller's mastiff bat, *Molossus pretiosus* (Mammalia: Chiroptera), from the Brazilian Caatinga. *Chiroptera Neotropical*, 14(1), 346–353.
- Novaes, R.L.M., & Laurindo, R.S. 2014. Morcegos da Chapada do Araripe, Nordeste do Brasil. *Papéis Avulsos de Zoologia*, 54 (22), 1-13. DOI: 10.1590/0031-1049.2014.54.22
- Nunes, H., Rocha, P.A., Sales, J., Rocha, F.L., & Cordeiro-Estrela, P. 2018. First record of *Uroderma magnirostrum* Davis, 1968 (Chiroptera: Phyllostomidae) in the northeastern Atlantic Forest of Brazil. *Oecologia Australis*, 22(3), 312-319. DOI: 10.4257/oeco.2018.2203.10
- Oliveira, E.R. 2000. Espaço ecomorfológico da fauna de quirópteros da região do Alto Tocantins, GO: uma perspectiva através da análise da morfologia alar. Master thesis. Universidade Federal do Rio de Janeiro. p. 82.
- Pacheco, S. M. 2004. Técnicas de campo empregadas no estudo de quirópteros. *Caderno La Salle* XI, 1(2), 193-205.
- Paglia, A.P., Fonseca, G.A.B., Rylands, A.B., Herrmann, G., Aguiar, L.M.S., Chiarello, A.G., Leite, Y.L.R., Costa, L.P., Siciliano, S., Kierulff, M.C.M., Mendes, S.L., Tavares, V.C., Mittermeier, R.A., Patton, J.L. 2012. Lista anotada dos mamíferos do Brasil. 2nd ed. Arlington: Occasional Papers in Conservation Biology, 6, 1-75.
- Papavero, N. 1994. Fundamentos práticos de taxonomia zoológica: coleções, bibliografia, nomenclatura. N. Papavero (Ed.), São Paulo, UNESP: p. 291.
- Passos, F.C., Silva, W.R., Wagner, A.P., & Bonin, M.R. 2003. Frugivoria em morcegos (Mammalia, Chiroptera) no Parque Estadual Intervales, sudeste do Brasil. *Revista Brasileira de Zoologia*, 20(3), 511–517. DOI: 10.1590/S0101-81752003000300024
- Pavan, A.C., Bobrowiec, P.E.D., & Percequillo, A.R. 2018. Geographic variation in a South American clade of mormoopid bats, *Pteronotus* (Phyllodia), with description of a new species. *Journal of Mammalogy*, 99(3), 624-645. DOI: 10.1093/jmammal/gyy048

- Peracchi, A.L., & Albuquerque, S.T. 1993. Quirópteros do Município de Linhares, Estado do Espírito Santo, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Biologia*, 53(1), 575-581.
- Pereira, M.J., Marques, J.T., Santana, J., Santos, C.D., Valsecchi, J., Queiroz, H.L., Beja, P., & Palmeirim J.M. 2009. Structuring of Amazonian bat assemblages: the roles of flooding patterns and floodwater nutrient load. *Journal of Animal Ecology*, 78(6), 1163-1171. DOI: 10.1111/j.1365-2656.2009.01591.x
- Peters, S.L., Malcolm, J.R., Zimmerman, B.L. 2006. Effects of Selective Logging on Bat Communities in the Southeastern Amazon. *Conservation Biology*, 20(5), 1410-1421. DOI:10.1111/j.1523-1739.2006.00526.x.
- Pina, S.M., Meyer, C.F., & Zortéa, M.A. 2013. Comparison of habitat use by phyllostomid bats (Chiroptera: Phyllostomidae) in natural forest fragments and Eucalyptus plantations in the Brazilian Cerrado. *Chiroptera Neotropical*, 19(3), 14-30.
- Pine, R.H., Bishop, I.R., & Jackson, R.L. 1970. Preliminary list of mammals of the Xantina/Cachimbo expedition (Central Brazil). *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 64(5), 668-670. DOI:10.1016/0035-9203(70)90003-9.
- Pinto, S.R.L. 2008. Inferências filogeográficas e estruturação populacional de *Sturnira lilium* (Phyllostomidae) da Mata Atlântica. Master thesis. Universidade Federal do Espírito Santo. p. 56.
- Pires, D.P.S., & Cademartori, C.V. 2018. Influence of the size and degree of conservation of remnants of the Atlantic rain forest on chiropteran assemblages. *Acta Biológica Catarinense*, 5(2), 68-79. DOI: 10.21726/abc.v5i2.595
- Presley, S.J., Willig, M.R., Wunderle, J.J.M., & Saldanha, L.N. 2008. Effects of reduced-impact logging and forest physiognomy on bat populations of lowland Amazonian forest. *Journal of Applied Ecology*, 45(1), 14–25. DOI: 10.1111/j.1365-2664.2007.01373.x

- Reis, N.M., Fregonezi, M.N., Peracchi, A.L., & Shibatta, A.O. 2013. Morcegos do Brasil: guia de campo. Rio de Janeiro: Technical Books: p. 252.
- Rocha, P.A., Soares, F.A., Dias, D., Mikalauskas, J.S., Anderson, F., Vilar, E.M., & Daher, M.R. 2017. New records of *Micronycteris schmidtorum* Sanborn, 1935 (Phyllostomidae, Chiroptera) for northeastern Brazil. *Mastozoología neotropical*, 24(2), 475-482.
- Sampaio, E., Lim, B. & Peters, S. 2016. *Sturnira tildae*. The IUCN Red List of Threatened Species 2016. e.T20960A22050501. (Retrieved on August 15th, 2019, from <http://dx.doi.org/10.2305/IUCN.UK.2016-2.RLTS.T20960A22050501.en>).
- Santos, F.C.B., Lisboa, C.V., Xavier, S.C.C., Dario, M.A., Verde, R.S., Calouro, A.M., Roque, A.L.R., & Jansen, A.M. 2017. *Trypanosoma* sp. diversity in Amazonian bats (Chiroptera; Mammalia) from Acre State, Brazil. *Parasitology*, 145(6), 828-837. DOI:10.1017/S0031182017001834
- Santos, T.C.M. 2017. Diversidade de morcegos em unidades de conservação na Amazônia. Master thesis. Instituto Tecnológico Vale. p. 84.
- Silva, R.C. 2016. Efeitos da estratificação vertical e obstrução vegetacional sobre a assembleia de morcegos de dois fragmentos florestais do leste do estado do Acre, Brasil. Master thesis. Universidade Federal do Acre. p. 61.
- Silva, R.M.T., & Bobrowiec, P.E.D. 2015. Diagnóstico da fauna de morcegos em unidades de conservação estaduais na área de influência direta da BR-319, estado do Amazonas. In: M. Gordo & H.S. Pereira (Eds.), *Unidades de Conservação do Amazonas no interflúvio Purus-Madeira: Diagnóstico Biológico*. Manaus, EDUA: pp. 193-206.
- Simmons, N.B. 2005. Order Chiroptera. In: D.E. Wilson & D.M. Reeder (Eds.), *Mammals Species of the World: a taxonomic and geographic reference*. Baltimore, Johns Hopkins University Press: pp. 312-529.

- Simmons, N.B., & Voss, R.S. 1998. The mammals of Paracou, French Guiana: a Neotropical lowland rainforest fauna, part 1, Bats. *Bulletin of the American Museum of Natural History*, 237(1), 1-219.
- Souza, R.F., Novaes, R.L.M., Siqueira, A., Sauwen, C., Jacob, G., Santos, C.E.L., Felix, S., Ribeiro, E., Sant'Anna, C., Vrcibradic, D., Avilla, L., Sbragia, I., & Santori, R.T. 2015. Morcegos (Mammalia, Chiroptera) em remanescente de Floresta Atlântica, Rio de Janeiro, sudeste do Brasil. *Neotropical Biology and Conservation*, 10(1), 9-14. DOI:10.4013/nbc.2015.101.02
- Taddei, V.A. 1975. Phyllostomidae (Chiroptera) do norte ocidental do Estado de São Paulo. II – Glossophaginae; Carollinae; Sturnirinae. *Ciência & Cultura*, 27(1), 723-734.
- Timm, R.M. 1994. The mammal's fauna. In: L.A. Mcdade, K.S. Bawa, H.A. Hespeneide, & G.S. Hartshorn (Eds.), *La Selva: ecology and natural history of a neotropical rainforest*. Illinois, University of Chicago Press: pp. 229-237.
- Velazco, P.M., & Patterson, B.D. 2013. Diversification of the yellow-shouldered bats, genus *Sturnira* (Chiroptera, Phyllostomidae), in the New World tropics. *Molecular Phylogenetics and Evolution*, 68(3), 683-698. DOI: 10.1016/j.ympev.2013.04.016
- Velazco, P. & Patterson, B.D. 2017. *Sturnira lilium*. The IUCN Red List of Threatened Species 2017: e.T88159688A22049384. (Retrieved on August 15th, 2019, from <http://dx.doi.org/10.2305/IUCN.UK.2017-2.RLTS.T88159688A22049384.en>).
- Velazco, P.M., & Patterson B.D. 2019. Small Mammals of the Mayo River Basin in Northern Peru, with the description of a new species of *Sturnira* (Chiroptera: Phyllostomidae). *Bulletin of the American Museum of Natural History*, 429, 1-70.
- Velazco, P.M., Gardner, A.L., & Patterson, B.D. 2010. Systematics of the *Platyrrhinus helleri* species complex (Chiroptera: Phyllostomidae), with descriptions of two new species. *Zoological Journal of the Linnean Society*, 159(3), 785–812. DOI:10.1111/j.1096-3642.2009.00610.x

Velazco, P.M., Soto-Centeno, J.A., Fleck, D.W., Voss, R.S., & Simmons, N.B. 2017. A new species of nectar-feeding bat of the genus *Hsunitycteris* (Phyllostomidae: Lonchophyllinae) from northeastern Peru. *American Museum Novitates*, 3881, 1–26. DOI: 10.1206/3881.1

Submitted: 22 May 2019

Accepted: 22 October 2019

Associate Editor: Alexandra Bezerra

Ahead of print