

*Macrobrachium potiuna* preying on adult treefrog

**STOMACH MORPHOLOGY AND FIRST RECORD OF THE  
FRESHWATER SHRIMP *MACROBRACHIUM POTIUNA* PREYING ON  
ADULT TREEFROG IN THE ATLANTIC FOREST**

Thiago Novato<sup>1\*</sup>, Arthur Diesel Abegg<sup>2</sup>, Bruno Ferreto Fiorillo<sup>3,4</sup>

<sup>1</sup> Universidade Federal Rural de Pernambuco, Programa de Pós-Graduação em Etnobiologia e Conservação da Natureza, Laboratório de Sociobiodiversidade, Rua Dom Manuel de Medeiros, s/n, Dois Irmãos, CEP: 52171-900, Recife, PE, Brazil.

<sup>2</sup> Instituto Butantan, Laboratório Especial de Coleções Zoológicas, Av. Vital Brasil, 1500, Butantã, CEP: 05503-900, São Paulo, SP, Brazil.

<sup>3</sup> Herp Trips, Rodovia SP-139, km 80, São Miguel Arcanjo, SP, Brazil.

<sup>4</sup> Instituto Manacá, Rodovia SP-139, km 80, Zona Rural, s/n, Abaitinga, CEP: 18230-000, São Miguel Arcanjo, SP, Brazil.

E-mail: thiago.novato@icb.ufjf.br (\*corresponding author); arthur\_abegg@hotmail.com; brunoferreto@alumni.usp.br

**Abstract:** We present the first report of *Macrobrachium potiuna* (Arthropoda: Decapoda) preying on an adult anuran, alongside with a description of the stomach morphology of the shrimp individual. The freshwater shrimp was observed feeding on a juvenile hylid in the Atlantic Forest. We interpret the shrimp's behavior as opportunistic predation instead of necrophagy due to the small size of the treefrog and the absence of decomposition. We also briefly discuss some ecological theories that might be used to explain this trophic relationship. Shrimp dissection revealed no gastric mill or other maceration structures, but the stomach contents included dendrites and tissues of the prey, as well as two insect eggs from the order Diptera. Our findings contribute to the limited knowledge on the diet, anatomy, and trophic relationships of freshwater shrimps.

**Keywords:** Amphibia; Southeastern Brazil; Crustacea; Predation

---


Freshwater shrimps are crucial for maintaining the trophic structure of lotic environments as they have an omnivorous diet that promotes nutrient cycling and energy flow in headwater streams (Carvalho *et al.* 2011, Cruz *et al.* 2021). They feed on organic plant debris in suspension or in the substratum (Carvalho *et al.* 2011, Jimoh *et al.* 2011, Lima *et al.* 2014), as well as on insects, oligochaetes, microcrustaceans, mollusks, and small fish (Mirzajani *et al.* 2020). For example, the predation of dipteran larvae by the shrimp *Macrobrachium inpa* Kensley & Walker, 1982 can regulate the population dynamics of these insects (Landeiro *et al.* 2008). This feeding pattern is also observed in the predation of ephemeropterans by *Macrobrachium americanum* Bate, 1868 (Moulton *et al.* 2004).

Although less frequent, there are reports of anuran eggs and tadpoles as food items for some freshwater shrimp and crab species (Warkentin 1999, Warren *et al.* 2021). Other arthropods can prey on juvenile forms of anurans, such as spiders (Nyffeler & Altig 2020), harvestmen (Menegucci *et al.* 2020), and water bugs (González-Maya *et al.* 2019). The predation risk for tropical anurans is higher during their reproductive period peak, which for most tropical species is occurs in the wet and warm season. During this period (between October and March in southeastern Brazil), higher anuran activity can increase their exposure to predators (Duellman & Trueb 1994). However, the frequency of post-metamorphic anuran predation rates by invertebrates is highly dependent on the presence of specialized tactics by the predators (Toledo *et al.* 2007). For example, smaller invertebrates with predatory tactics such as venom or social foraging behavior are more likely to consume prey of equal or larger size, while non-venomous or solitary invertebrates tend to consume smaller prey (Toledo *et al.* 2007).

On December 10, 2021 (summer), during a fieldwork carried out at the Reserva Particular do Patrimônio Natural Trápaga, São Miguel Arcanjo municipality, in the state of São Paulo, southeastern

Brazil (24°3'20.938"S; 47°58'48.414"W, WGS84; 784 m a.s.l.), we observed a male individual of *Macrobrachium potiuna* Müller 1880 (carapace length of 45 mm) feeding on a small-sized juvenile treefrog (snout-vent length ca. 25 mm, ~ 55% of the shrimp size) (Figure 1). These anurans are primarily arboreal and usually inhabit stream beds, bromeliads, or rocky cavities in the Atlantic Forest (Frost 2011). Both shrimp and treefrog individuals were collected under ICMBio and SISBIO permits (#73355, #73380, #79423-2, and #79423-3) and deposited at the Didactic Entomological Collection of the Zoology Department of the Universidade Federal de Juiz de Fora.


Ahead of print



**Figure 1.** *Macrobrachium potiuna* feeding on adult treefrog at the Reserva Particular do Patrimônio Natural Trápaga, São Miguel Arcanjo, São Paulo, Brazil (A); collected individual of *M. potiuna* (B). Photo credits: Bruno Ferreto Fiorillo.

*Macrobrachium potiuna* “*stricto sensu*” is distributed across upland forest areas in the states of Santa Catarina, Paraná, and São Paulo, occurring in sandy streams with diverse marginal vegetation (Müller & Carpes 1991, Mattos & Oshiro 2009). Although studies have been conducted on its reproductive biology, population dynamics, and taxonomy (Mattos & Oshiro 2009; Pileggi & Mantelatto 2013), no study to date reported on its diet or foraging strategies. However, information on the diet composition allows to identify feeding habits, determine the trophic position, and distinguish species (Lima *et al.* 2014, Mirzajani *et al.* 2020). Furthermore, the description of a species' digestive system (dimensions and shapes) can provide insights into its habits and feeding preferences (Karasov *et al.* 2011). To enhance our knowledge regarding the trophic relationships of *M. potiuna* in natural systems and to provide new natural history information on the species, we described the morphology of the digestive tract and the food items found within it, following the methods proposed by Lima *et al.* (2016).

The *M. potiuna* individual had a 3 mm long stomach (Figure 2), and its cardiac stomach was located posterior to the head and lacked a gastric mill. The cardiac sac was concave and had rough, striated muscles. During the dissection process, the stomach and pyloric filter were accidentally dismembered. Inside the stomach, we identified tissues from a hylid species, along with detritus and two Diptera eggs.



**Figure 2.** Esophagus and stomach morphology of *Macrobrachium potiuna*. CS = Cardiac stomach; Cfil = Cardiac filament; CSac = Cardiac sac; Eso = Esophagus. Illustration credits: Dra. Samyra Furtado.

We presume that the observed predation event was opportunistic because there are no similar cases reported for other species of *Macrobrachium*, despite the predatory habit of other freshwater shrimp species (Warren *et al.* 2021). We discard the possibility of necrophagy because usually the eye socket is the first structure to decompose, and the prey eye socket did not show any signals of decomposition. In addition, the shrimp individual was twice as large as the anuran, indicating the possibility of predation.

We indicate two potential hypotheses to explain this behavioral episode. The first is related to changes in food availability for *M. potiuna*. December 2021 was characterized by intense rainfall concentrated in a few hours of the day (Agritempo 2022). Such intense rainfall increases the water flow and the current in streams that, despite stirring up the suspended benthic substrate, can also

reduce the availability of the preferred food for *Macrobrachium* (Esteves 2011), such as eggs and larval stages of insects (Minshall & Minshall 1977, Bunn & Arthington 2002). The second hypothesis is related to the increasing foraging activity that occurs during the reproductive peak of shrimps *Macrobrachium* spp. in December (Sampaio *et al.* 2007). Obtaining body biomass and energy are essential for the production and maturation of viable sperm in males, in addition to success in their competition for females (Pérez-Rodríguez *et al.* 2019).

According to Carvalho *et al.* (2013), the populations of *M. potiuna* split into two different lineages, “*stricto sensu*”, distributed across the Brazilian Plateau, and “*Affinis-Clade*”, distributed in floodplain areas of the states of Minas Gerais and Bahia. In this context, research that investigates morphological differences in the digestive tract and diet among different clades is desirable, since environmental conditions (apart from biogeographic factors) can influence the availability of food resources (Tundisi & Tundisi 2008) that can potentially lead to morphological differences amongst clades (Goodheart *et al.* 2017).

As far as we know, this is the first report of adult anuran predation by *Macrobrachium* and the first available diet record of *M. potiuna*. Moreover, we present the first description of its digestive tract morphology. We highlight that amphibians are important in the composition of food webs (Blaustein *et al.* 2011, Davenport & Chalcraft 2012, Zipkin *et al.* 2020), and emphasize the urgent need for quantitative research on predator-prey relationships between crustaceans and amphibians that can deepen the knowledge of the basic ecology and trophic levels of these groups.

## REFERENCES

- AgriTempo. Retrieved on February 19, 2022, from <https://www.agritempo.gov.br/agritempo/index.jsp>
- Blaustein, A. R., Han, B. A., Relyea, R. A., Johnson, P. T. J., Buck, J. C., Gervasi, S. S., & Kats, L. B. 2011. The complexity of amphibian population declines: Understanding the role of cofactors in driving amphibian losses. *Annals of the New York Academy of Sciences*, 1223(1), 108–

119. DOI: 10.1111/j.1749-6632.2010.05909.x

Bunn, S. E., & Arthington, A. H. 2002. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental Management*, 30(4), 492–507. DOI: 10.1007/s00267-002-2737-0

Carvalho, D. A., Collins, P. A., & De Bonis, C. J. 2011. Gut evacuation time of *Macrobrachium borellii* (Caridea: Palaemonidae) feeding on three types of prey from the littoral-benthic community. *Journal of Crustacean Biology*, 31(4), 630–634. DOI:10.1651/10-3446.1

Carvalho, F. L., Pileggi, L. G., & Mantelatto, F. L. 2013. Molecular data raise the possibility of cryptic species in the Brazilian endemic prawn *Macrobrachium potiuna* (Decapoda, Palaemonidae). *Latin American Journal of Aquatic Research*, 41(4), 707–717. DOI:10.3856/vol41-issue4-fulltext-7

Cruz, B. R. F., Cunha, M. C., Bueno, A. A. D. P., & Jacobucci, G. B. 2021. Natural diet of *Macrobrachium brasiliense* (Crustacea, decapoda) in a cerrado stream. *Iheringia - Série Zoologia*, 111, 1–11. DOI: 10.1590/1678-4766e2021003

Esteves, F. A. 2011. *Fundamentos de Limnologia*. Editora Interciência, Rio de Janeiro, Brasil: 826 p.

Goodheart, J.A., Bazinet, A.L., Valdés, Á., Collins, A. G., & Cummings, M. P. 2017. Prey preference follows phylogeny: evolutionary dietary patterns within the marine gastropod group *Cladobranchia* (Gastropoda: Heterobranchia: Nudibranchia). *BMC Ecology and Evolution*, 17(1), DOI: 10.1186/s12862-017-1066-0

Davenport, J. M., & Chalcraft, D. R. 2012. Evaluating the effects of trophic complexity on a keystone predator by disassembling a partial intraguild predation food web. *Journal of Animal Ecology*, 81(1), DOI:10.1111/j.1365-2656.2011.01906.x

Duellman, W. E., & Trueb, L. 1994. *Biology of amphibians*. Maryland: The Johns Hopkins University Press: p 696.

Frost D.R. 2011. *Amphibian species of the world: An online reference*. Version 5.5 (31 January 2011): <http://research.amnh.org/vz/herpetology/amphibia/>, American Museum of Natural



History, New York, USA.

- González-Maya, J. F., Gómez-Hoyos, D. A., Seisedos-De-vergara, R., Cruz-Lizano, I., & Schipper, J. 2019. Water-bug (*Abedus* sp.; Belostomatidae) predation on the critically endangered *Atelopus Varius* (bufonidae) at las tablas protected Zone, Costa Rica. *Acta Biologica Colombiana*, 24(2), 403–406. DOI:10.15446/abc.v24n2.76924
- Jimoh, A. A., Clarke, E. O., Whenu, O. O., & Adeoye, H. B. 2011. Food and feeding habits of the African river prawn (*Macrobrachium vollenhovenii*, Herklots, 1857) in Epe Lagoon, southwest Nigeria. *International Journal of Fisheries and Aquaculture*, 3(1), 10–15.
- Karasov, W. H., Martínez, R. C., & Caviedes-Vidal, E. 2011. Ecological physiology of diet and digestive systems. *Annual Review of Physiology*, 73(1), 69–93. DOI: 10.1146/annurev-physiol-012110-142152
- Landeiro, V. L., Hamada, N., & Melo, A. S. 2008. Responses of aquatic invertebrate assemblages and leaf breakdown to macroconsumer exclusion in Amazonian “terra firme” streams. *Fundamental and Applied Limnology*, 172(1), 49–58. DOI: 10.1127/1863-9135/2008/0172-0049
- Lima, J. F., Garcia, J. S., & Silva, T. C. 2014. Natural diet and feeding habits of a freshwater prawn (*Macrobrachium carcinus*: Crustacea, Decapoda) in the estuary of the Amazon River. *Acta Amazonica*, 44(2), 235–244. DOI: 10.1590/s0044-59672014000200009
- Lima, J. F., Garcia, J. S., & Tavares, M. 2016. Foregut morphology of *Macrobrachium carcinus* (Crustacea, Decapoda, Palaemonidae). *Acta Amazonica*, 46(2), 209 – 218. DOI: 10.1590/1809-4392201501214
- Mattos, L. A., & Oshiro, L. M. Y. 2009. Population structure of *Macrobrachium potiuna* (Crustacea, Palaemonidae) in the Moinho’s River, Mangaratiba, Rio de Janeiro, Brazil. *Biota Neotropica*, 9(1), 81–86. DOI: 10.1590/S1676-06032009000100010
- Melo, M. S., & Nakagaki, J. M. 2013. Evaluation of the feeding habits of *Macrobrachium brasiliense*

- (Heller, 1862) in the Cural de Arame stream (Dourados/Mato Grosso Do Sul, Brazil).  
Nauplius, 21(1), 25–33. DOI: 10.1590/s0104-64972013000100004
- Menegucci, R. C., Muscat, E., Tanaka, R. M., & Stuginski, D. 2020. Predation on *Hylodes phyllodes* (Anura: Hylodidae) by the harvestman *Heteromitobates discolor* (Arachnida: Opiliones).  
Phyllomedusa, 19(1), 121–124. DOI: 10.11606/issn.2316-9079.v19i1p121-124
- Minshall, G. W., & Minshall, J. N. 1977. Microdistribution of benthic invertebrates in a rocky mountain (U.S.A.) Stream. Hydrobiologia, 55(1), 231–249.
- Mirzajani, A., Ghane, A., Bagheri, S., Abbasi, K., Sayadrahim, M., Salahi, M., & Lavajoo, F. 2020. Diet survey and trophic position of *Macrobrachium nipponense* in the food web of Anzali wetland. Wetlands, 40(5), 1229–1239. DOI: 10.1007/s13157-020-01278-5
- Moulton, T. P., De Souza, M. L., Silveira, R. M. L., & Krsulović, F. A. M. 2004. Effects of ephemeropterans and shrimps on periphyton and sediments in a coastal stream (Atlantic forest, Rio de Janeiro, Brazil). Journal of the North American Benthological Society, 23(4), 868–881. DOI: 10.1899/0887-3593(2004)023<0868:EOEASO>2.0.CO;2
- Müller, Y. M. R., & Carpes, S. 1991. *Macrobrachium potiuna* (Müller): aspectos do ciclo reprodutivo e sua relação com parâmetros ambientais (Crustacea, Decapoda, Palaemonidae). Revista Brasileira de Zoologia, 8(1–4), 23–30. DOI: 10.1590/s0101-81751991000100004
- Nyffeler, M., & Altig, R. 2020. Spiders as frog-eaters: A global perspective. Journal of Arachnology, 48(1), 26–42. DOI: 10.1636/0161-8202-48.1.26
- Pérez-Rodríguez, J. C., Gómez-Gutiérrez, J., López-Greco, L. S., & Cortés-Jacinto, E. 2019. Spermatophore production and sperm quality of the river prawn *Macrobrachium americanum* Spence Bate, 1868 fed with different diets. Aquaculture Research, 50(11), 3117–3129. DOI: 10.1111/are.14265
- Pileggi, L. G., & Mantelatto, F. L. 2013. Taxonomic revision of doubtful Brazilian freshwater shrimps species of genus *Macrobrachium* (Decapoda, Palaemonidae). Iheringia. Série Zoologia,

102(4), 426–437. DOI: 10.1590/s0073-47212012005000012

Sampaio, C. M. S., Silva, R. R., Santos, J. A., & Sales, S. P. 2007. Reproductive cycle of *Macrobrachium amazonicum* females (Crustacea, Palaemonidae). Brazilian Journal of Biology, 67(3), 551–559. DOI: 10.1590/s1519-69842007000300022

Toledo, L. F., Ribeiro, R. S., & Haddad, C. F. B. 2007. Anurans as prey: an exploratory analysis and size relationships between predators and their prey. Journal of Zoology, 271(2), 170–177. DOI: 10.1111/j.1469-7998.2006.00195.x

Tundisi, J.G., & Matsumura-Tundisi, T. 2008. Limnologia. São Paulo: Oficina de Textos, 632. Wetzel, R.G. Limnología. Barcelona: Ediciones Omega S.A., 1981.679 p.

Warkentin, K. M. 1999. The development of behavioral defenses: a mechanistic analysis of vulnerability in red-eyed tree frog hatchlings. Behavioral Ecology, 10 (3), 251-262. DOI: 10.1093/beheco/10.3.251.

Warren, D. A., Bradbeer, S. J., & Dunn, A. M. 2021. Superior predatory ability and abundance predicts potential ecological impact towards early-stage anurans by invasive ‘Killer Shrimps’ (*Dikerogammarus villosus*). Scientific Reports, 11(1), 1–12. DOI: 10.1038/s41598-021-82630-5

Zipkin, E. F., DiRenzo, G. V.; Ray, J. M., Rossman, S., & Lips K. R. 2020. Tropical snake diversity collapses after widespread amphibian loss. Science 367(6479), 814–816. DOI: 10.1126/science.aay573

*Submitted: 01 August 2022*

*Accepted: 28 August 2023*

*Published online: 11 September 2023*

*Associate Editor: Bruno Soares*

Ahead of print