



ICHTHYOFAUNA OF THE MARIMBONDINHO RIVER, FURNAS RESERVOIR, MINAS GERAIS, BRAZIL

André Batista Nobile^{1}, Valter Monteiro Azevedo-Santos^{2,3,4}, Claudio Lopes Soares⁵,
Felipe Viana Manzano⁵, Claudio Oliveira⁶ & Felipe Pontieri de Lima¹*

¹ Ictiológica Consultoria Ambiental LTDA-ME, Rua Primo Paganini, 80, CEP 18608-190, Botucatu, SP, Brazil.

² Faculdade Eduvale de Avaré, Avenida Prof. Misael Eufrásio Leal, Centro, CEP: 18705-050, Avaré, São Paulo, Brazil

³ Universidade Federal do Tocantins, Programa de Pós-Graduação em Biodiversidade, Ecologia e Conservação, Rua 03, Quadra 17, Lote 11, S/N, CEP: 77500-000, Porto Nacional, Tocantins, Brazil.

⁴ Espaço Inovação do Parque de Ciência e Tecnologia Guamá, Grupo de Ecologia Aquática, Rua da Inovação, S/N, Belém, Pará, Brazil.

⁵ FURNAS Centrais Elétricas S.A., Av. Graça Aranha, 26, Rio de Janeiro, RJ, CEP 20030-000 Brazil.

⁶ Universidade Estadual Paulista, Instituto de Biociências, Departamento de Biologia Estrutural e Funcional, R. Prof. Dr. Antônio Celso Wagner Zanin, 250, CEP 18618-689, Botucatu, SP, Brazil.

E-mails: andrenobile@hotmail.com (*corresponding author); valter.ecologia@gmail.com; clsoares@furnas.com.br; fmanzano@furnas.com.br; claudio.oliveira@unesp.br; fpl.limao@hotmail.com.

Abstract: Ichthyofauna surveys are essential for conservation efforts in reservoirs. Here we present a list of fish species found in the Marimbondinho River, a tributary of the Furnas HPP reservoir in Minas Gerais, Brazil. Fish captures were conducted using gillnets and cast nets at five sampling points during six campaigns in 2018 and 2019. We recorded a total of 1,892 individuals belonging to 31 species, distributed in 16 families and four orders. Four fish species were considered long-distance migrants, and five were identified as non-natives. None of the captured species are listed as threatened on the official Brazilian list. Order Characiformes exhibited the highest species richness; in which Anostomidae and Characidae were the families with great species richness. *Hoplosternum littorale* was the most abundant species, presented in approximately 22% of all the captures. Overall, the results indicate that the ichthyofauna of Marimbondinho River is similar to those typically observed in Brazilian reservoirs, with the expected richness and number of migratory species. We expect this study can help in future monitoring programs and, fundamentally, to propose conservation and management measures on endemic and migratory species of Furnas reservoir.

Keywords: anthropic impacts; ichthyofauna survey; fish ecology; Neotropical fishes.

The change from a lotic to lentic environment by the construction of a hydroelectric plant causes drastic variations in the ichthyofauna (Agostinho *et al.* 2008). For instance, Furnas Hydro Power Plant (Rio Grande, Upper Paraná River) is among the oldest reservoirs in Southeast Brazil — and has impacted fish assemblages since its construction. Studies on the ichthyofauna of this reservoir have focused, for example, on the aquaculture (Furnas 2007), biology of fish species (Santos *et al.* 2004, Ribeiro *et al.* 2007), and introduction of non-native species (Azevedo-Santos *et al.* 2011). However, many regions of the Furnas reservoir, especially in the mouths of tributaries, have no information on fish species or assemblages; this is the case, for example, of the Marimbondinho

River. A survey on the ichthyofauna of this region can contribute in different ways, as in addition to generating information that supports new studies, it helps in the identification of migratory species that use the region to complete their reproductive cycle. Therefore, here we provide an ichthyofauna survey of the Marimbondinho River, a tributary located upstream of the Furnas reservoir, in the municipality of Boa Esperança, Minas Gerais, Brazil. This study is part of what was requested by the Public Prosecutor's Office of the District of Boa Esperança – Minas Gerais, in the records of the Civil Inquiry (IC) nº MPMG – MPMG-0071.19.000163-7.

The Furnas reservoir is one of the largest in Brazil, with 1,440 km² of flooded area (Azevedo-Santos *et al.* 2011). Dammed in 1963, it flooded part of its main rivers, such as the Grande and Sapucaí, and many tributaries, such as the Marimbondinho River, which flows into the upper part of the Grande River, near the municipality of Boa Esperança, Minas Gerais (21°5'48.94" S, 45°33'21.98" W). Many cities bordering the Furnas reservoir make use of it for tourism, and this is the case of Boa Esperança municipality. Due to issues related to the large variation in the level of the reservoir that directly affected the region of the mouth of the Marimbondinho River and brought sanitary problems to the city of Boa Esperança, a dike was built in the middle course of the river, close to the city, giving rise to Lago dos Encantos. The presence of this dike, in addition to reducing the effects of the discharge of sewage from the city into its dammed waters, regulates the level of the dam upstream and allows tourism activities throughout the year in the city. In this way, the construction of the dike reduced the influence of the reservoir of the Furnas HPP in the upstream

areas, while the downstream areas are susceptible to annual variation in its level. Thus, the sampling program was defined in order to sample five sites (S1 to S5) (Table 1; Figure 1), two located upstream of the dike, in Lago dos Encantos (S1 and S2), and three downstream from it, in the flooded arm of the reservoir of the Furnas HPP itself (S3 to S5).

Six samplings were carried out at each station (ABIO authorization No. 1099/2019), in the months of May, August, November and December of 2019 and January and February of 2020. The samplings were carried out using a set of eight gill nets (meshes of 3, 5, 6, 7, 8, 10, 12, 14 cm, between opposite nodes for 25 meters in length), which were installed perpendicular to the bank, at dusk, and removed the following day (approximately 14 hours of exposure). Due to fluctuations in the level of the reservoir at Furnas HPP, the complete set was installed in all campaigns only at points S1, S2 and S5 (Supplementary Material 1). Cast nets (diameter of 3 m and meshes of 2 to 3 cm) were also used with a total of 15 castings of netting at each point.

The captured individuals were placed in plastic bags containing data on the location and mesh size. After, fish were identified using keys (Menezes 1987, Oyakawa & Mattox 2009, Ota *et al.* 2018, Peixoto *et al.* 2015), counted, photographed, weighed and the total length (TL) and standard (SL) measured. Representative specimens (*i.e.*, vouchers) of species were fixed in formaldehyde (10%) and preserved in alcohol (70%) and deposited in the collection of the Laboratory of Fish Biology (LBP), at the Universidade Estadual Paulista "Júlio de Mesquita Filho", in Botucatu, São Paulo, Brazil.

The species were classified according to their origin and their migratory behavior based on the

Table 1. Characterization of each sampling site used in this study to perform ichthyofauna surveys along Marimbondinho River, Minas Gerais, Brazil.

Station	Coordinates	Position	Substrate	Riparian vegetation	Depth (m)
S1	21° 4'17.78"S; 45°33'43.73"W	Upstream	Sand	Absent	5
S2	21° 4'54.01"S; 45°31'46.43"W	Upstream	Sand	Absent	5
S3	21° 3'56.36"S; 45°33'13.45"W	Downstream	Organic matter	Macrophytes bank	1,5
S4	21° 2'59.80"S; 45°32'57.04"W	Downstream	Organic matter	Macrophytes bank	2,5
S5	21° 1'37.74"S; 45°32'50.52"W	Downstream	Organic matter	Macrophytes bank	4




Figure 1. Location of sampling points along the Marimbondinho, Boa Esperança, Minas Gerais, Brazil (modified from Azevedo-Santos *et al.* 2019).

literature (Langeani *et al.* 2007, Vazzoler 1996, Agostinho *et al.* 2003, Vasconcelos *et al.* 2014, Santos 2010). The classification of families into orders, and species into families, followed the Eschmeyer's Catalog of Fishes (Fricke *et al.* 2023, Van der Laan *et al.* 2023). The species rarefaction curve was estimated, considering all samples taken at all points as replicates. The observed richness (Sobs) and Bootstrap were used as estimators. These analyses were performed using Primer 6.0 software (Anderson *et al.* 2008).

A total of 1,892 individuals was captured, representing 31 species, 16 families and four

orders (Supplementary Material 2). Four species were classified as long-distance migratory, *Prochilodus lineatus* (Valenciennes 1837) (N = 6), *Salminus hilarii* Valenciennes 1850 (N = 1), *Megaleporinus obtusidens* (Valenciennes 1837) (N = 6) and *Pimelodus maculatus* Lacepède 1803 (N = 97) (Supplementary Material 2). Five species were classified as non-native, three allochthonous (*Cichla piquiti* Kullander & Ferreira 2006, *Hoplosternum littorale* (Hancock 1828) and *Metynniss lippincottianus* (Cope 1870)), and two exotics (*Coptodon rendalli* (Boulenger 1897) and *Oreochromis niloticus* (Linnaeus 1758)).

Characiformes was the order with greater richness and abundance, with 18 species (58% of the total) and 1,026 individuals (54.2%), followed by Siluriformes, with seven species (22% of the total) and 728 individuals (38.5%) (Figure 2). Among the 16 families, the highest richness was observed in Anostomidae, with five species, followed by Characidae and Cichlidae, both with four species (Figure 2). In terms of abundance, Characidae was the most important family, with 526 individuals (27.8%), followed by Callichthyidae (22.2%) and Curimatidae (19.3%). *Hoplosternum littorale* was the most abundant, with 419 individuals (22.1%), followed by *Steindachnerina insculpta* (Fernández-Yépez 1948) (18.5%) and *Psalidodon fasciatus* (Cuvier 1819) (15.9%) (Figure 3).

The highest abundance was found in S5, which was captured 680 individuals, followed by S1, with 547 (Figure 4). The site with the highest species richness was S2, with 22 species, followed by S5, with 21 (Figure 5). The species accumulation curve (Figure 6) approached the plateau, according to the methods used (Sobs = 31; Bootstrap = 32.72).

The data presented here indicate that the ichthyofaunistic richness of the sampled sites is lower than expected for dammed environments, where between 40 and 50 species were found by

other authors (e.g., Pelicice *et al.* 2018, Nobile *et al.* 2019b, Queiroz-Sousa *et al.* 2019). However, it is important to point out that the use of gillnets may have exerted sample selectivity on the ichthyofauna, preventing, for instance, the capture of small-sized species, such as those of the Poeciliidae and Characidae family.

Regarding the four migratory species recorded, it was observed that except for *Salminus hilarii*, which had only one individual captured upstream of the dike at S2, the others had greater abundance recorded at sites downstream of the dike. The Furnas HPP reservoir receives several tributaries, from streams to rivers, the main tributaries being the Grande and Sapucaí rivers, which have upstream stretches that have not yet been dammed. Migratory fish need dam-free stretches, as they do not reproduce in lentic ecosystems, which is one of the main reasons why dams represent disturbances to migratory populations (Marques *et al.* 2018, Agostinho *et al.* 2008). The capture of these long-distance migrants in the upper reaches of the Furnas HPP reservoir indicates that its tributaries can play an important role in maintaining these populations and, in this sense, the preservation of the natural flow of these waterbodies becomes crucial. However, this does not seem to be the




Figure 2. Abundance and richness by orders and families recorded along the Marimbondinho stream, Boa Esperança, Minas Gerais, Brazil. line = abundance; columns = richness.




Figure 3. Number of individuals recorded for the most abundant species along the Marimbondinho, Boa Esperança, Minas Gerais, Brazil.

case in the Marimbondinho River region, due to the occurrence of few migratory species and the low abundance recorded for these, with no schools being recorded even in the spawning months, only the presence of a few individuals.

Five non-native species were recorded. *Hoplosternum littorale* is attributed as a native of the upper Paraná system (Langeani *et al.* 2007). However, Santos (2010) reported that the species was introduced into the Rio Grande basin. *Cichla piquiti* probably occurs in the studied area due to sport fishing activities, and fishing tournaments still occur in neighboring municipalities. On the other hand, the presence of *C. rendalli* and *O. niloticus* in the reservoir is consequence of the aquaculture activity, since these species were widely introduced in Brazil by this activity (Nobile *et al.* 2019a), including in the Furnas HPP reservoir

(Azevedo-Santos *et al.* 2011). The occurrence of *M. lippincottianus* in the region may be result of aquarium dumping, since this is one explanation to its introduction in other regions of the country (Assis *et al.* 2017).

The order Characiformes dominated in terms of richness and abundance, followed by the order Siluriformes (Figure 2). In the Neotropical region as a whole, whether in reservoirs or in environments under lower anthropic pressure, this pattern has been treated as expected, with some inversion between these orders (Lowe-McConnell 1987, Agostinho & Júlio Jr 1999, Langeani *et al.* 2007, Jarduli *et al.* 2020).

In this study, the highest species richness per family was recorded for Anostomidae. This constitutes a peculiarity, since, generally, Characidae and Loricariidae are the families with

the greatest richness (Langeani *et al.* 2007, Jarduli *et al.* 2020, Pelicice *et al.* 2018). These results may also be linked to the selectivity of the capture methods, and, in this study, the low richness observed for the Characidae family corroborates with some studies

that used only gillnets (Britto & Carvalho 2006, Nobile *et al.* 2019b). As anostomids generally reach medium to large sizes, they may have been captured more frequently with the methods employed, contributing to greater richness.




Figure 4. Fish abundance per sampling sites along the Marimbondinho River arm, Boa Esperança, Minas Gerais, Brazil.




Figure 5. Fish richness per sampling sites along the Marimbondinho River arm, Boa Esperança, Minas Gerais, Brazil.

Despite the lower richness, the highest abundances per family were recorded for Characidae, Curimatidae and Callichthyidae. In environments under the effect of dams, such as those sampled in this study, changes in the composition and structure of assemblages may occur, generally with a decrease in large species, more common in lotic environments, and an increase in small ones (Agostinho *et al.* 2016, Lima *et al.* 2016, Nobile *et al.* 2019b). In this scenario, in the upstream environments (S1 and S2), Characidae, represented by *A. lacustris* and *P. fasciatus*, and Curimatidae, represented by *S. insculpta*, express a population explosion, since those species adjusted very well to these new environments (Ribeiro *et al.* 2007, Peressin *et al.* 2012, Lopes *et al.* 2016). The third most abundant family, Callichthyidae, was more representative at the downstream points (S3 to S5) (Figure 2), represented almost exclusively by *H. littorale*, the most abundant species in this study (22% of catches) (Figure 3). *Hoplosternum littorale* exhibits parental care (Vazzoler 1996); builds nests (Ramnarine 1995) and before and after hatching they protect the offspring against predators, which can explain its success. Also, they do not depend

on running water to reproduce (Vazzoler 1996), feed on various types of items (Vazzoler *et al.* 1997, Souto *et al.* 2016) and tolerate environments with less oxygenation (Brauner *et al.* 1995), as that observed downstream.

In this study, *Oligosarcus argenteus* Günther 1864 was a new record for the Furnas reservoir. Azevedo-Santos *et al.* (2019), studying streams, provided the report of its occurrence for the Paraná River basin. This species probably occurs naturally both in streams and in the reservoir at Furnas HPP. Thus, this record is an expansion in the distribution of *O. argenteus* in the Paraná basin.

Eigenmannia cf. trilineata (López & Castello 1966) is considered a species very similar to other congeners with subterminal mouth, as previously described (Peixoto *et al.* 2015, Peixoto & Wosiacki 2016, Campos-da-Paz & Queiroz 2017). Apparently, the species we captured in the region is the same one reported by Azevedo-Santos *et al.* (2019) from an affluent creek of the reservoir of Furnas HPP. The exact identification of the species will depend on new morphological analyses (and involving osteological characteristics). With these new assessments, it will be possible to determine




Figure 6. Species accumulation curve considering all samplings and collection stations in the Marimbondinho River, Boa Esperança, Minas Gerais, Brazil.

whether this is a species already described (*e.g.*, *E. besouro* Peixoto & Wosiacki 2016 or if it is a possibly undescribed species of the “*E. trilineata* group” (*sensu* Peixoto *et al.* 2015).

The ichthyofauna of the Marimbondinho River was highly impacted by humans, with reduced species diversity, with low abundance of migrants, but with species adapted to lentic and/or dammed environments, as well as the presence of non-native species. We hope that our study will provide a baseline to help future monitoring programs in Furnas reservoir; and, therefore, help to establish appropriate conservation and management measures — especially focusing on endemic and migratory species.

ACKNOWLEDGEMENTS

We are grateful to Ictiológica Consultoria Ambiental for all support during the collection. To Furnas Centrais Elétricas S.A. for giving data obtained in the “Estudo da dinâmica ecológica e populacional da ictiofauna e análise da viabilidade de implantação de mecanismo de transposição de peixes no dique do Lago dos Encantos, Boa Esperança (MG), no reservatório da UHE Furnas” and allow us the publication of the results. We wish to thank two anonymous reviewers, the editors, and Donald C. Taphorn B. for important suggestions. CO received financial support from Fundação de Amparo à Pesquisa do Estado de São Paulo - FAPESP grants 2018/20610-1, 2016/09204-6, 2014/26508-3 and Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq proc. 306054/2006-0.

REFERENCES

- Agostinho, A. A., & Júlio Jr, H. F. 1999. Peixes da bacia do Alto Paraná. In: R. H. Lowe-McConnell (Ed.), Estudos Ecológicos de Comunidades de Peixes Tropicais. pp. 374–400. São Paulo: Edusp.
- Agostinho, A. A., Pelicice, F. M., & Gomes, L. C. 2008. Dams and the fish fauna of the Neotropical region: impacts and management related to diversity and fisheries. *Brazilian Journal of Biology*, 68(4), 1119–1132. DOI: 10.1590/S1519-69842008000500019
- Agostinho, A. A., Gomes, L. C., Santos, N. C. L., Ortega, J. C. G., & Pelicice, F. M. 2016. Fish assemblages in Neotropical reservoirs: Colonization patterns, impacts and management. *Fisheries Research*, 173, 26–36. DOI: 10.1016/j.fishres.2015.04.006
- Agostinho, A. A., Gomes, L. C., Suzuki, H. I., & Júlio Jr, H. F. 2003. Migratory fishes of the Upper Paraná River Basin, Brazil. In: B. Carolsfeld, C. R. Harvey, & A. Baer (Eds.), *Migratory fishes of South America: biology, fisheries and conservation status*. pp. 19–98. Ottawa: World Fisheries Trust, International Bank for Reconstruction and Development, The World Bank.
- Anderson, M. J., Gorley, R. N. ., & Clarke, K. R. 2008. *Permanova + for Primer: guide to software and statistical methods*. Plymouth: PRIMER-E.
- Assis, D. A. S. de, Dias-Filho, V. A., Magalhães, A. L. B., & Brito, M. F. G. 2017. Establishment of the non-native fish *Metynnis lippincottianus* (Cope 1870) (Characiformes: Serrasalminidae) in lower São Francisco River, northeastern Brazil. *Studies on Neotropical Fauna and Environment*, 52(3), 228–238. DOI: 10.1080/01650521.2017.1348057
- Azevedo-Santos, V. M., Rigolin-Sá, O., & Pelicice, F. M. 2011. Growing, losing or introducing? Cage aquaculture as a vector for the introduction of non-native fish in Furnas Reservoir, Minas Gerais, Brazil. *Neotropical Ichthyology*, 9(4), 915–919. DOI: 10.1590/S1679-62252011000400024
- Azevedo-Santos, V. M., Britski, H. A., Oliveira, C., & Benine, R. C. 2019. Ichthyofauna of streams of the Rio Sapucaí basin, upper Rio Paraná system, Minas Gerais, Brazil. *Biota Neotropica*, 19(1). DOI: 10.1590/1676-0611-bn-2018-0617
- Brauner, C. J., Ballantyne, C. L., Randall, D. J., & Val, A. L. 1995. Air breathing in the armoured catfish (*Hoplosternum littorale*) as an adaptation to hypoxic, acidic, and hydrogen sulphide rich waters. *Canadian Journal of Zoology*, 73(4), 739–744. DOI: 10.1139/z95-086
- Britto, S. G. C., & Carvalho, E. D. 2006. Ecological attributes of fish fauna the Taquaruçu reservoir, Paranapanema River (Upper Paraná, Brazil): composition and spatial distribution. *Acta Limnologica Brasiliensia*, 18, 377–388.
- Campos-da-Paz, R., & Queiroz, I. R. 2017. A new species of *Eigenmannia* Jordan and Evermann (Gymnotiformes: Sternopygidae) from the upper rio Paraguai basin. *Zootaxa*, 4216(1), 73.

- DOI: 10.11646/zootaxa.4216.1.5
- Fricke, R., Eschmeyer, W. N., & Van der Laan, R. 2023. Eschmeyer's Catalog Of Fishes: Genera, Species, References. Retrieved on July 18, 2023, from <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>
- Furnas. 2007. Estudo Técnico-Científico Visando a Delimitação de Parques Aquícolas nos Lagos das Usinas Hidroelétricas de Furnas e Três Marias – MG. Relatório de Identificação de Áreas Tecnicamente Adequadas para a Instalação de Parques Aquícolas. Reservatório de Furna. p. 348.
- Jarduli, L. R., Garcia, D. A. Z., Vidotto-Magnoni, A. P., Casimiro, A. C. R., Vianna, N. C., Almeida, F. S. de, Jerep, F. C., & Orsi, M. L. 2020. Fish fauna from the Paranapanema River basin, Brazil. *Biota Neotropica*, 20(1). DOI: 10.1590/1676-0611-bn-2018-0707
- Langeani, F., Castro, R. M. C., Oyakawa, O. T., Shibatta, O. A., Pavanelli, C. S., & Casatti, L. 2007. Diversidade da ictiofauna do Alto Rio Paraná: composição atual e perspectivas futuras. *Biota Neotropica*, 7(3), 181–197. DOI: 10.1590/S1676-06032007000300020
- Lima, A. C., Agostinho, C. S., Sayanda, D., Pelicice, F. M., Soares, A. M. V. M., & Monaghan, K. A. 2016. The rise and fall of fish diversity in a neotropical river after impoundment. *Hydrobiologia*, 763(1), 207–221. DOI: 10.1007/s10750-015-2377-z
- Lopes, V. G., Nessimian, J. L., Da-Silva, E. R., Gomes, J. H. C., Dias, A. C. I. M., Souza, L. C., & Branco, C. W. C. 2016. Habitat heterogeneity on feeding habit of two sympatric and congeneric characidae fishes in two tropical reservoirs. *Iheringia. Série Zoologia*, 106. DOI: 10.1590/1678-4766e2016012
- Lowe-McConnell, R. H. 1987. Ecological studies in tropical fish communities. Cambridge: Cambridge University Press: p. 391.
- Marques, H., Dias, J. H. P., Perbiche-Neves, G., Kashiwaqui, E. A. L., & Ramos, I. P. 2018. Importance of dam-free tributaries for conserving fish biodiversity in Neotropical reservoirs. *Biological Conservation*, 224, 347–354. DOI: 10.1016/j.biocon.2018.05.027
- Menezes, N. A. 1987. Três espécies novas de *Oligosarcus* Gunther, 1864 e redefinição taxonômica das demais espécies do gênero (Osteichthyes, Teleostei, Characidae). *Boletim de Zoologia*, 11(11), 1. DOI: 10.11606/issn.2526-3358.bolzoo.1987.122368
- Nobile, A. B., Cunico, A. M., Vitule, J. R. S., Queiroz, J., Vidotto-Magnoni, A. P., Garcia, D. A. Z., Orsi, M. L., Lima, F. P., Acosta, A. A., da Silva, R. J., do Prado, F. D., Porto-Foresti, F., Brandão, H., Foresti, F., Oliveira, C., & Ramos, I. P. 2019a. Status and recommendations for sustainable freshwater aquaculture in Brazil. *Reviews in Aquaculture*, 12(3), 1495–1517. DOI: 10.1111/raq.12393
- Nobile, A. B., Freitas-Souza, D., Lima, F. P., Queiroz, J., Bayona-Perez, I. L., Carvalho, E. D., & Ramos, I. P. 2019b. Damming and seasonality as modulators of fish community structure in a small tributary. *Ecology of Freshwater Fish*, 28(4), 563–572. DOI: 10.1111/eff.12475
- Ota, R. R., Deprá, G. de C., Graça, W. J. da, & Pavanelli, C. S. 2018. Peixes da planície de inundação do alto rio Paraná e áreas adjacentes: revised, annotated and updated. *Neotropical Ichthyology*, 16(2). DOI: 10.1590/1982-0224-20170094
- Oyakawa, O. T., & Mattox, G. M. T. 2009. Revision of the Neotropical trahiras of the *Hoplias lacerdae* species-group (Ostariophysi: Characiformes: Erythrinidae) with descriptions of two new species. *Neotropical Ichthyology*, 7(2), 117–140. DOI: 10.1590/S1679-62252009000200001
- Peixoto, L. A. W., & Wosiacki, W. B. 2016. *Eigenmannia besouro*, a new species of the *Eigenmannia trilineata* species-group (Gymnotiformes: Sternopygidae) from the rio São Francisco basin, northeastern Brazil. *Zootaxa*, 4126(2), 262. DOI: 10.11646/zootaxa.4126.2.6
- Peixoto, L. A. W., Dutra, G. M., & Wosiacki, W. B. 2015. The Electric Glass Knifefishes of the *Eigenmannia trilineata* species-group (Gymnotiformes: Sternopygidae): monophyly and description of seven new species. *Zoological Journal of the Linnean Society*, 175(2), 384–414. DOI: 10.1111/zoj.12274
- Pelicice, F. M., Azevedo-Santos, V. M., Esguícero, A. L. H., Agostinho, A. A., & Arcifa, M. S. 2018. Fish diversity in the cascade of reservoirs along the Paranapanema River, southeast Brazil. *Neotropical Ichthyology*, 16(2). DOI: 10.1590/1982-0224-20170150
- Peressin, A., Gonçalves, C. da S., & Braga, F. M. de S. 2012. Reproductive strategies of

- two Curimatidae species in a Mogi Guaçu impoundment, upper Paraná River basin, São Paulo, Brazil. *Neotropical Ichthyology*, 10, 847–854. DOI: 10.1590/S1679-62252012000400018
- Queiroz-Sousa, J., Keith, S. A., David, G. S., Brandão, H., Nobile, A. B., Paes, J. V. K., Souto, A. C., Lima, F. P., Silva, R. J., Henry, R., & Richardson, K. 2019. Species richness and functional structure of fish assemblages in three freshwater habitats: effects of environmental factors and management. *Journal of Fish Biology*, 95(4), 1125–1136. DOI: 10.1111/jfb.14109
- Ramnarine, I. W. 1995. Induction of nest building and spawning in *Hoplosternum littorale*. *Journal of Fish Biology*, 47(3), 555–557. DOI: 10.1111/j.1095-8649.1995.tb01921.x
- Ribeiro, V. M. A., Santos, G. B., & Bazzoli, N. 2007. Reproductive biology of *Steindachnerina inculpta* (Fernandez-Yépez) (Teleostei, Curimatidae) in Furnas reservoir, Minas Gerais, Brazil. *Revista Brasileira de Zoologia*, 24(1), 71–76. DOI: 10.1590/S0101-81752007000100009
- Santos, J. E. dos, Bazzoli, N., Rizzo, E., & Santos, G. B. 2004. Reproduction of the catfish *Iheringichthys labrosus* (Lütken) (Pisces, Siluriformes) in Furnas reservoir, Minas Gerais, Brazil. *Revista Brasileira de Zoologia*, 21(2), 193–200. DOI: 10.1590/S0101-81752004000200006
- Santos, G. B. 2010. A ictiofauna da bacia do Alto Paraná (rio Grande e rio Paranaíba). *MG. Biota*, 2(6), 20.
- Souto, A. C., Lima, F. P., Gildo, N. M., & Vidotto-Magnoni, A. P. 2016. Dieta das Espécies de Peixes do Reservatório de Jurumirim e seus Principais Tributários. In: R. J. da Silva (Ed.), *Integridade ambiental da represa de Jurumirim: Ictiofauna e relações ecológicas*. São Paulo: Editora Unesp: p. 335.
- Van der Laan, R., Fricke, R., & Eschmeyer, W. N. 2023. *Eschmeyer's catalog of fishes: Classification*.
- Vasconcelos, L. P., Alves, D. C., & Gomes, L. C. 2014. Fish reproductive guilds downstream of dams. *Journal of Fish Biology*, 85(5), 1489–1506. DOI: 10.1111/jfb.12501
- Vazzoler, A. E. A. M. 1996. *Biologia da Reprodução de peixes Teleósteos: Teoria e Prática*. Maringá: EDUEM: p. 169.
- Vazzoler, A. E. A. M., Lizama, M. A. P., & Inada, P. 1997. Influências ambientais sobre a sazonalidade reprodutiva. In: A. E. A. M. Vazzoler, A. A. Agostinho, & N. S. Hahn (Eds.),

A planície de inundação do alto rio Paraná: aspectos físicos, biológicos e socioeconômicos. pp. 267–280. Maringá: Eduem.

SUPPLEMENTARY MATERIAL

Supplementary Material 1. Effort applied at each sampling station, in different campaigns.

Supplementary Material 2. Taxa (orders, families, and species) and abundance (in each sampling station) found in the Furnas HPP reservoir, Boa Esperança region, Minas Gerais, southeastern Brazil. * non-natives; ** migrators.

Submitted: 6 June 2023

Accepted: 1 September 2023

Published online: 11 September 2023

Associate Editor: Felipe Ottoni