



ICHTHYOFAUNA FROM AN AMAZONIAN ISLAND: DIVERSITY, REPRODUCTION, AND FEEDING ASPECTS

Danielly Torres Hashiguti¹, Antonio Augusto Jardim Junior², Élide Jesana Santana Cunha², Thaize dos Santos Souza¹ & Thierry Frédou³*

¹ Universidade Federal do Pará, Núcleo de Ecologia Aquática e Pesca da Amazônia, R. do INPE, S/N - Terra Firme, CEP 66077-530, Belém, PA, Brazil.

² Universidade Federal do Pará, Instituto de Ciências Biológicas, Laboratório de Ecologia e Conservação, Av. Perimetral, 2-224 - Guamá, CEP 66077-830, Belém, PA, Brazil.

³ Universidade Federal Rural do Pernambuco, Departamento de Pesca e Agricultura, R. Manuel de Medeiros, S/N - Dois Irmãos, CEP 52171-900, Recife, PE, Brazil.

Email: danigtorres@gmail.com (*corresponding author); biosantosjr@gmail.com; elidajesana@gmail.com; thaize_souza@yahoo.com.br; tfredou@depaq.ufrpe.br.

Abstract: The high productivity in estuaries supports a high biomass of fishes, making them important environments for feeding, development, and reproduction of larvae and juveniles of many economically and ecologically important species. Many islands compose the landscape of the Amazon estuary given the large deposition of soil caused by the Amazon River waterflow. Given the importance of estuarine environments to coastal ecosystems, this study aims to diagnose the fish community inhabiting a typical island of the Amazon estuary and describe the temporal patterns in community composition, reproductive and feeding activity. Specimens were collected in the main channel and a tidal creek of Onças Island, quarterly between September 2006 and 2008, using gillnets and block-off nets. Captured fishes were identified, sexed, measured (total length and weight), had reproductive and feeding activities described, and were classified into environmental and feeding guilds. Abundance and biomass were expressed by the Capture per Unit of Effort in number of individuals (CPUEn) and biomass (CPUEb). The island was characterized as a nursery ground (high abundance of individuals with immature and at maturity gonads), reproduction ground (high abundance of individuals with mature and spawned/spent gonads) and feeding ground (high abundance of individuals with stomachs containing food). A total of 37 species distributed in 21 families and 1,177 individuals were captured. Richness varied significantly throughout the collections but there were no significant differences in fish abundance (CPUEn) or biomass (CPUEb) between periods in any habitat type. Individuals displayed different feeding activity between hydrological periods and feeding guilds. Freshwater species dominated through all collection periods and most species were zoobenthivore, piscivore, or planktivore. Our results indicated that the island serves as a feeding and reproduction ground for several freshwater fish species, possibly reflecting the importance of other islands in Amazonian estuarine systems and highlighting their importance for conservation.

Keywords: Feeding; Guilds; List of species; Nursery; Onças Island.

INTRODUCTION

Estuaries are transition zones between marine and freshwater ecosystems where the salinity is constantly changing (Dame 2008). On average, estuarine waters are biologically more productive than those of rivers and the adjacent ocean due to hydrodynamic characteristics of the circulation, which stimulates productivity by trapping nutrients, algae, and plants on the system (Chilton *et al.* 2021). Due to greater productivity and high environmental variability, estuaries support high abundance and biomass of fishes, generally with dominance of few species adapted to varying salinities (Rundle *et al.* 1998; Camargo *et al.* 2003; Castro & Huber 2012), including both secondary freshwater species and primary marine species (Barletta & Lima 2019).

Seasonal changes in the river discharge constantly alter environmental conditions, including salinity, which is a major variable shaping biological communities in coastal areas (Barletta *et al.* 2005; Dantas *et al.* 2010; Vergès *et al.* 2022; Soares *et al.* 2021; Marceniuk *et al.* 2023). For example, freshwater species may use estuaries during the entire year, but they become more abundant during the rainy season when there is increased discharge of continental waters into the estuary. The opposite happens in the dry season, when sea water invades estuaries and primarily marine fishes become more frequent (Potter *et al.* 2010). Therefore, the ichthyofauna that inhabits estuaries as feeding, breeding, and nursery grounds are a combination of freshwater, estuarine resident, and marine species (Selleslagh *et al.* 2009; Pichler *et al.* 2015; Molina *et al.* 2020), of which many are economically and ecologically important (Chaves *et al.* 2002; Franco *et al.* 2008; Guebert-Bartholo *et al.* 2011).

In addition to having high ecological value, estuaries suffer constant anthropogenic pressure (Dias *et al.* 2013, De Araújo-Barbosa *et al.* 2016), which is one of the leading factors in the decline of estuarine biodiversity and habitat quality, leading to the loss of ecosystem services worldwide (Firth *et al.* 2016, Chaudhary *et al.* 2018). Among the multiple anthropogenic impacts that affect estuarine fishes are: (i) overfishing, which contributes to the reduction of fish populations and unbalance of community dynamics (Raz-Guzmán & Huidobro 2002); (ii) biological invasions that interfere with local trophic relationships (Barletta & Lima 2019); (iii)

discharges from industrial pollutants, agricultural waste and domestic sewage, damaging water quality and directly threatening the fishes' health (Costa *et al.* 2012); (iv) climate change, including rising water temperatures, ocean acidification, and sea level rise, impacting environmental conditions and the capacity of fishes to deal with seasonal changes (Blaber 2013); and (v) the loss or change of critical habitats, compromising essential areas for reproduction and feeding (Viana *et al.* 2010).

These direct human disturbances also threaten the environmental conditions of the Amazon estuary, the largest in terms of freshwater discharge in the world (Kennish 2002; Latrubesse *et al.* 2017; Lessa *et al.* 2018). The enormous discharge carries considerable amounts of nutrients and organic matter and creates a very unstable coastal region, subject to erosion processes that result in high deposition of soil that contributes to the development of a large number of islands (Lessa *et al.* 2018). Surrounding Belém, one of the largest Brazilian cities in the Amazon that is home for more than 1,3 M inhabitants (IBGE 2022), extends a set of Amazonian estuarine islands (Gregório & Mendes, 2009) that are inhabited by multiple euryhaline fishes of high commercial value for industrial fisheries and the subsistence of riverside human communities (Bentes *et al.* 2012; Viana & Frédou 2014; Marceniuk *et al.* 2021a; Prestes *et al.* 2022). Since 1980, the region has been subject to the effects of ore and soy outflow through a large port project (CDP 2023), but only in the last decade a rampant tourism started to develop (Nunes & Furtado 2023), and the lack of scientific information about these stressors and its effects in the aquatic environment hamper the development of mitigation and conservation strategies.

Given the importance of understanding how species use different ecosystems and their responses to environmental changes (Beck *et al.* 2003), this study aims to analyze the fish community inhabiting an Amazonian island under anthropogenic activities, using a dataset prior to the recent intensification of the described impacts. We tested the hypothesis that the island served as a nursery area, feeding, and breeding ground for fish species, and we expect to generate a baseline to understand how the establishment and intensification of threats will affect the local ichthyofauna.

MATERIAL AND METHODS

Study area

The Onças Island, it is approximately 19 km long, and is part of the Barcarena County in the state of Pará and is at 2.5 km northwest of the capital Belém by waterway (). The climate is tropical type Af in the Köppen system (Peel *et al.* 2007), and during the collection period the accumulated values of monthly rainfall in the capital Belém showed a pattern similar to the mean climate based on the average of the last 30 years (INMET 2013), with well-defined rainy (March), rainy/dry (June), dry (September), and dry/rainy (December) periods.

Water temperature (28.6 ± 0.65 °C), pH (6 ± 0.45), and salinity (0 ppm) do not show significant variation throughout the year. Tidal variation is the most striking environmental factor in the region, reaching up to 3.6 m in spring tide conditions (Gregório & Mendes 2009) and creating multiple tidal channels that cross the island. Tidal channels, hereafter named as tidal creeks, are narrow inlets near marine coastlines exhibiting water fluctuations and currents according with the tide regimen, and are very common in estuarine islands (Healy, 2005).

Sampling and Laboratorial Procedures

At the tidal creek, sampling was carried out quarterly between September 2006 and 2008 in nights of spring tide using a block nylon multifilament net (35 m x 7 m and 0.75 mm knot-to-knot mesh) set at the mouth of a creek (Figure 1, Site number 2). At the river channel, sampling was carried out quarterly between September 2007 and 2008 in nights of spring tide using monofilament nylon gill nets (knot-to-knot meshes of 25 mm, 30 mm, 40 mm and total length of 210 m) set in three different points at the river channel (Figure 1, Site number 1). Different knot-to-knot mesh sizes were selected for each environment because small fishes tend to avoid open waters and nets with small knot-to-knot mesh sizes could be damaged and/or obstructed with the high volume of debris in the main channel. The block net was fixed to the bottom of the creek during the low tide and raised during high tide while buoyancy devices allowed them to occupy the entire water column when the water levels increased, then

capturing all individuals that entered the habitat for approximately 12 hours. The main channel was sampled twice per sampling expedition, with nets being underwater for nine hours each, between 6 am and 6 pm.

Individuals were captured under the SISBio collection license number 17250-1 and identified according to the identification guide of Van Der Sleen and Albert (2017) and with the help of experts for each taxonomic group. In the laboratory, specimens had their total length (TL; 0.1 cm) and total weight (TM; 0.1 g) measured. One ventral-longitudinal incision was performed to remove the gonads and stomachs. The gonadal development was classified using a macroscopic scale adapted from Núñez and Duponchelle (2009): A – immature (Females: small, circular, thin, and pinkish ovaries with no oocytes visible to the naked eye; Males: small and silvery/translucent testes); B – maturing (Female: bigger ovaries filled with white or yellowish oocytes of different sizes; Male: testes are longer and with whitish to pinkish color); C – mature or ripe (Female: the ovary reaches its maximal development and the oocytes can be expelled with a light pressure; Male: testes are fuller, completely white, and sperm can be expelled with a light pressure); and D – spent (Female: ovaries are empty, often bloody, with ripe oocytes still visible; Male: large, flaccid and empty testes). The macroscopical characterization of gonadal stages is proven effective for several fish species and the estimation of reproductive parameters such as the length at first maturity (Hashiguti *et al.* 2017; 2019; Soares *et al.* 2020), especially for community-level approaches and studies with budget and logistic limitations.

For feeding aspects, our objective was exclusively to assess whether individuals were feeding in the main channel and/or accessing tidal creeks to feed, and the stomachs were categorized according to the repletion degree as “with food” (WF) and “no food” (NF, the absence of organic material in stomach content). Although stomach content analysis provides a narrow view of recent food items, it is still a common and widely accepted method for categorization (Amundsen and Sánchez-Hernández, 2019).




Figure 1. Sampling stations in the (1) Main Channel and (2) Tidal Creek in Onças Island, Pará, Brazil.

Data analysis

Species diversity was characterized by the number of species (observed richness), the equity in the distribution of individuals among species (Pielou's evenness, J), abundance, and biomass. Using each sampling event in main channel and tidal creek as samples, observed richness was tested using a Kruskal-Wallis test (H) with 5% significance level, followed by Dunn's multiple comparisons test (Dunn 1961) for significative results. Considering the differences in sample size and sampling effort between habitat type (tidal creek and main channel), we chose to use an individual-based (based on the abundance of the community standardized by the smallest number of individuals) rarefaction method with the iNEXT package (iNterpolation and EXTrapolation) in the R program (R Core Team 2022), using the Hill number estimator for species richness ($q=0$) (Hill 1973), to compare communities between habitat types. The Capture per Unit of Effort in number of individuals (CPUEn) and biomass (CPUEb) were calculated using the following equations: $CPUE_n = (n/A * t) * 100$ and $CPUE_b = (b/A * t) * 100$ where A = the net area (m^2) and t = the immersion time (hours). Differences in CPUEn and CPUEb between grouped hydrological periods (rainy, rainy/dry, dry, and dry/rainy) for each habitat type (main channel and tidal creek) were tested by Analysis of Variance (ANOVA), followed by a Tukey post-hoc for significative results (Zar 1996).

The habitat type was classified as a reproduction site for the species that displayed significant amounts of individuals with mature (C) and spent (D) gonads; and as feeding area characterized by the predominance of individuals with stomachs with food (WF). The frequency of feeding (WF vs. NF) and reproducing (A/B vs. C/D) fish was compared using chi-squared tests across hydrological periods and habitat type. The effect of the trophic group over feeding individuals was evaluated using the percentage of stomachs with organic content (WF) for each feeding group using an ANOVA. Species were classified in feeding (DV: detritivore; OV: omnivore; PL: planktivore; PV: piscivore; CV: carnivore; HV: herbivore and ZB: zoobenthivore) and environmental guilds (ES: estuarine; FS: freshwater; FM: freshwater migrants) according to Franco *et al.* (2008) and Fishbase (Froese & Pauly 2018). Differences in feeding guilds (abundance) by

collection periods (rainy, rainy/dry, dry, and dry/rainy) and habitat type (main channel and tidal creek) were tested by a PERMANOVA using a Bray Curtis similarity matrix (Zar 1996). All analyzes were carried out using the R program (R Core Team 2022) using the "vegan" package (Oksanen *et al.* 2019), with a significance level of 5%.

RESULTS

A total of 1,177 individuals belonging to 37 species and 21 families were captured (Table 1 and Appendix A), with 25 species captured at the main channel (9 exclusive) and 27 captured at the tidal creek (12 exclusive). The Aspredinidae, Sciaenidae and Pimelodidae families correspond to 65% of all collected individuals. Evenness ($J = 0.7 \pm 0.1$) was similar throughout collection period, but observed richness was significantly higher in March 2008 (Appendix A; $H = 18.02$; $p = 0.0009$). The rarefied richness was similar between habitat types (overlapping confidence intervals), with an estimate of 39 species for the island with an increase effort (main channel = 29 and tidal creek = 30; Appendix B). There were no significant differences in the fish abundance (CPUEn) or biomass (CPUEb) between periods in any habitat type (Figure 2).

In terms of abundance and biomass, the most representative species in the main channel and tidal creek were similar, in except for *Lithodoras dorsalis* and *Aspredo aspredo*, which were one of the most representative ones in tidal creek and main channel, respectively (Figure 3). In general, the species caught were small and medium-sized or juveniles of larger species, with approximately 80% specimens with less than 25 cm total length (Appendix C), with a predominance of females (67%) and immature individuals ($A = 64\%$). The creek harbored a higher number of immatures and feeding individuals than the main channel, as well as larger and heavier specimens (Table 2). The island serves as reproduction area for 19 species, functioning both as nursery and reproduction grounds.

Among the 1,149 stomachs analyzed, approximately half presented stomach content (WF = 59%). Individuals fed differently throughout hydrological periods ($\chi^2 = 24$; $p < 0.001$) with higher amount of feeding individuals in September 2007 (dry period); habitat type ($\chi^2 = 5.40$; $p = 0.02$), with more feeding specimens in the tidal creek; and

Table 1. Species captured, range in total length (TL), feeding species (WF), gonadal development (A/B: Immature and maturing, C/D: mature and spent) feeding (PV: Piscivore; OV: Omnivore; HV: Herbivore, PL: Planktivore; ZB: Zoobenthivore, DV: Detritivore) and environmental guilds (ES: Estuarine, FS: Freshwater, FM: Freshwater Migrants), and place of occurrence on Onças Island, Pará, Brazil.

Order, Family and Species	TL	WF	Gonadal development		Feeding Guild	Environmental Guild	Main Channel	Tidal Creek	Total
			A/B	C/D					
ACANTHURIFORMES									
Sciaenidae									
<i>Micropogonias furnieri</i> (Desmarest, 1823)	15.5-24.5	46.67%	80%	20%	ZB	ES	15		15
<i>Pachypops fourcroyi</i> (Lacepède, 1802)	11.5-17.5	18.75%	25%	75%	ZB	FS	6	10	16
<i>Plagioscion squamosissimus</i> (Heckel, 1840)	13-30	75.85%	63.3%	36.7%	PV	FS	76	142	218
BELONIFORMES									
Belonidae									
<i>Strongylura timucu</i> (Walbaum, 1792)	34-34	0.0%	100%	0.0%	PV	ES	1		1
CHARACIFORMES									
Characidae									
<i>Psalidodon</i> aff. <i>fasciatus</i> (Cuvier, 1819)	6.7-11.5	25.0%	87.5%	12.5%	ZB	FS	1	11	12
Curimatidae									
<i>Curimata inornata</i> Vari, 1989	13-17.5	71.43%	100%	0.0%	HV	FS	7		7
Erythrinidae									
<i>Hoplias malabaricus</i> (Bloch, 1794)	17.4-20	33.33%	66.67%	33.33%	PV	FS		3	3
CICHLIFORMES									
Cichlidae									
<i>Cichla ocellaris</i> (Bloch & Schneider, 1801)	21-50	100%	50%	50%	PV	FS	1	1	2
<i>Crenicichla semifasciata</i> (Heckel, 1840)	23.5-36	33.33%	33.33%	66.67%	PV	FS	1	2	3
<i>Geophagus</i> gr. <i>altifrons</i> Heckel, 1840	14-19	0.0%	90%	10%	DV	FS	1	16	17
CLUPEIFORMES									
Engraulidae									
<i>Anchovia surinamensis</i> (Bleeker, 1865)	10-22	73.68%	45%	55%	PL	ES	2	18	20
<i>Lycengraulis batesii</i> (Günther, 1868)	8-24	22.62%	82.89%	17.11%	ZB	FM	46	39	85

Table 1. Continues on next page...

Table 1.continued

Order, Family and Species	TL	WF	Gonadal development		Feeding Guild	Environmental Guild	Main Channel	Tidal Creek	Total
			A/B	C/D					
Pristigasteridae									
<i>Pellona flavipinnis</i> (Valenciennes, 1837)	17-42.7	66.67%	100%	0.0%	PV	FM	6		6
GYMNOTIFORMES									
Gymnotidae									
<i>Gymnotus</i> gr. <i>carapo</i> Linnaeus, 1758	-	100%	100%	0.0%	PV	FS		1	1
Hypopomidae									
<i>Steatogenys elegans</i> (Steindachner, 1880)	22-22	0.0%	100%	0.0%	-	FS		4	4
Rhamphichthyidae									
<i>Rhamphichthys rostratus</i> (Linnaeus, 1766)	55-77	100%	60%	40%	ZB	FS		6	6
Sternopygidae									
<i>Sternopygus macrurus</i> (Bloch & Schneider, 1801)	38.5-39.7	100%	100%	0.0%	-	FS		2	2
<i>Sternopygus</i> sp.1	10.3-22.5	0.0%	100%	0.0%	-	FS		25	25
<i>Sternopygus</i> sp.2	32.4-46	87.5%	50%	50%	-	FS		8	8
PLEURONECTIFORMES									
Cyclopsettidae									
<i>Citharichthys spilopterus</i> Günther, 1862	19-19	33.33%	100%	0.0%	ZB	FM	2	2	4
SILURIFORMES									
Aspredinidae									
<i>Aspredo aspredo</i> (Linnaeus, 1758)	18-21	47.14%	84.62%	15.38%	ZB	ES	301		301
<i>Aspredinichthys filamentosus</i> (Valenciennes, 1840)	1-20.5	100%	50%	50%	ZB	ES	2		2
Auchenipteridae									
<i>Agenciosus</i> aff. <i>ucayalensis</i> Castelnau, 1855	13-27.5	63.54%	100%	0.0%	ZB	FS	14	82	96
<i>Trachelyopterus galeatus</i> (Linnaeus, 1766)	14.4-23.2	80%	100%	0.0%	OV	FS		5	5
Doradidae									
<i>Lithodoras dorsalis</i> (Valenciennes, 1840)	12-62.5	100%	100%	0.0%	OV	FS	2	45	47

Table 1. Continues on next page...

Table 1.continued

Order, Family and Species	TL	WF	Gonadal development		Feeding Guild	Environmental Guild	Main Channel	Tidal Creek	Total
			A/B	C/D					
Heptapteridae									
<i>Pimelodella cristata</i> (Müller & Troschel, 1849)	11.5-26.6	82.35%	63.33%	36.67%	ZB	FS		34	34
Loricariidae									
<i>Peckoltia</i> sp.1	11-13	0.0%	100%	0.0%	DV	FS	2		2
<i>Peckoltia</i> sp.2	13.5-13.5	0.0%	100%	0.0%	DV	FS		1	1
<i>Ancistrus</i> sp.1	10-12	25.0%	33.33%	66.67%	DV	FS	3	1	4
<i>Pseudacanthicus spinosus</i> (Castelnau, 1855)	13-13	0.0%	100%	0.0%	DV	FS	1		1
<i>Loricaria</i> cf. <i>cataphracta</i> Linnaeus, 1758	14-19	0.0%	100%	0.0%	DV	FS	1	3	4
Pimelodidae									
<i>Brachyplatystoma rousseauxii</i> (Castelnau, 1855)	17.5-17.5	100%	100%	0.0%	PV	FM	1		1
<i>Brachyplatystoma vaillantii</i> (Valenciennes, 1840)	32-38	100%	100%	0.0%	PV	FM		2	2
<i>Hypophthalmus marginatus</i> Valenciennes, 1840	13.5-44.5	35.44%	100%	0.0%	PL	FS	8	200	208
<i>Sorubim lima</i> (Bloch & Schneider, 1801)	28-34	100%	50%	50%	CV	FS	2		2
Pimelodidae									
<i>Pimelodus blochii</i> Valenciennes, 1840	16.5-27.5	100%	66.67%	33.33%	ZB	FS		9	9
TETRAODONTIFORMES									
Tetraodontidae									
<i>Colomesus asellus</i> (Müller & Troschel, 1848)	5.5-13	0.0%	33.33%	66.67%	OV	ES	2	1	3
Total Geral							504	673	1177




Figure 2. Capture per unit of effort in number (CPUEn) and biomass (CPUEb) by collection period (Rainy, transition from Rainy to Dry, Dry, and transition Dry to Rainy) and habitat type (Main channel and Tidal Creek), between September 2006 and September 2008, Onças Island, Pará, Brazil.

feeding guilds ($F = 11.1$; $p < 0.0001$), with carnivores ($WF = 100\%$), omnivores ($WF = 92.73\%$), piscivores ($WF = 74.9\%$), and herbivores ($WF = 71.43\%$) presenting more than half of individuals with stomach contents (Appendix D).

The abundance of environmental guilds was similar over hydrological periods ($F = 0.16$; $p = 0.96$) but significantly different when comparing habitat type ($F = 15.6$; $r^2 = 0.6$; $p = 0.002$). Freshwater species (FS) were dominant through all collection periods in the tidal creek while estuarine species (ES) were dominant in the main channel (Figure 4). Most species were zoobenthivorous ($ZB = 52\%$), piscivorous ($PV = 20\%$), or planktivorous ($PL = 19\%$) in the Onças Island (Figure 5). The abundance of feeding guilds differed between habitat types ($F = 7.03$; $r^2 = 0.41$ $p = 0.007$), but not through the year ($F = 0.642$; $r^2 = 0.19$ $p = 0.893$), with greater abundance

of zoobenthivorous species in the main channel and zoobenthivorous, piscivorous, and planktivorous species well distributed in the tidal creek (Figure 6).

DISCUSSION

The Onças island is strongly influenced by the Guajará Bay, featuring an environment influenced by continental waters and zero salinity, that served as a nursery, reproduction ground, and feeding ground for fishes, with individuals feeding differently throughout hydrological periods and feeding guilds. Richness varied significantly between periods, but there were no differences in the fish abundance (CPUEn) or biomass (CPUEb) between periods in any habitat type. Freshwater species dominated through all collection periods and most species were zoobenthivore, piscivore, or planktivore.




Figure 3. Percentage of the species with greater numerical and biomass participation in the main channel (a and b), and tidal creek (c and d), collected between September 2006 and September 2008, Onças Island, Pará, Brazil.

Table 2. Distribution of the ichthyofauna sex (M – Male, F- Female), Total Length in cm (TL), Total Weight in grams (TW), gonadal development (A - immature, B - at maturity, C – mature, and D – spawned or spent) and feeding behavior (NF - No Food, WF - With Food) among environments at Onças Island, Pará, Brazil.

Environment	Sex	TL (SD)	TW (SD)	Gonadal Development (%)				Feeding (%)	
				A	B	C	D	NF	WF
Tidal Creek	F = 74.34	22.9 ±9.87	102 ±156.88	42.86	17.86	38.1	1.19	34.51	65.49
	M = 25.66	22.1 ±8.67	112.1 ±248.9	57.47	27.59	14.94	0	28.74	71.26
	Total	23.18 ±9.09	109.81 ±199.69	71.7	10.73	17.11	0.47	42.84	57.16
Main Channel	F = 60.18	16.04 ±4.6	34.3 ±72.12	28.36	36.82	34.33	0.5	45.36	54.64
	M = 39.82	17.1 ±3.37	46.2 ±55.14	55.64	41.35	3.01	0	52	48
	Total	15.91 ±4.53	31.60 ±58.79	52.66	30.23	16.98	0.23	54.12	45.88
Tidal Creek and Main Channel	F = 67.36	19.9 ±8.54	72.1 ±130.09	36.42	26.27	36.42	0.88	39.2	60.8
	M = 32.64	18.9 ±6.36	72.2 ±165.44	56.36	35.91	7.73	0	42.45	57.55
	Total	19.89 ±8.22	76.32 ±160.50	64.03	18.55	17.05	0.37	40.24	59.76




Figure 4. Percentage of environmental guilds: (ES) estuarine species; (FS) freshwater species and (MM) marine migrants by dry (D), dry/rainy (D/R), rainy (R), and rainy/dry (R/D) periods, between September 2006 and September 2008, at Onças Island, Pará, Brazil.




Figure 5. Percentage of feeding guilds DV: detritivorous; OV: omnivorous; PL: planktivorous; PV: piscivorous; CV: carnivorous; HV: herbivorous and ZB: zoobenthivorous by dry (D), dry/rainy (D/R), rainy (R), and rainy/dry (R/D) periods, between September 2006 and September 2008, at Onças Island, Pará, Brazil.

The observed richness of Onças Island ($S = 37$) was low compared with estuaries receiving a stronger marine influence in the Amazon (73 species at the Curuçá River in Hercos 2006, 40 species at the Guajará Bay in Viana *et al.* 2010, Guajará and 108 species at the Marajó Bay and 94 at the Guajará

Bay in Mourão *et al.* 2015) and other regions (79 species at the Paraná Estuary in Barletta *et al.* 2008, 122 species at the Itapissuma, Suape, Sirinhaém, and Rio Formoso in Merigot *et al.* 2017, 102 species at the São Francisco River in Silva *et al.* 2023). No significant differences in the fish number and

biomass were observed over time, which could be a result of three factors. First, a very dynamic and highly disturbed surrounding area with strong tidal currents and wind generated waves (Pinheiro 2002, Ribeiro 2004) that may work as a geographic barrier. Second, the null salinity, since salinity is one of the most important environmental variable influencing fish assemblage and structure in estuaries (Bulger 1993, McLusky & Elliot 2004, Harrison & Whitfield 2006), where fish richness, abundance and biomass tend to increase with the rise in salinity at the lower estuary (predominance of marine species) and decrease with the river input at the upper estuary (predominance of freshwater species) (Whitfield & Harrison 2003, Kolpakov & Milovankin 2010, Whitfield *et al.* 2012). Third, sampling limitations, since we did not use other fishing gears (*e.g.* longlines and trawl nets) and sampled only on a few sites in the island (*e.g.* more tidal creeks and river channels). By doing so, abundant species in the region, such as *Cynoscion acoupa*, *Menticirrhus americanus*, *Plagioscion surinamensis*, *Stellifer microps* and *S. rastrifer* (Mourão *et al.* 2014), could possibly be registered.

The predominance of females among the captured species corresponds to the expected tendency of females to remain in estuarine systems (Secor 1999, Bartulović *et al.* 2004, Walsh *et al.* 2004, Callihan *et al.* 2013, Earl 2014), and the presence of 64% of immature individuals captured supports the idea of these areas as natural nurseries, important habitats for the life cycle of many fish species, providing protection against predators and food availability (Beck *et al.* 2001, Barletta *et al.* 2010, Barros *et al.* 2011). When comparing environments, the tidal creek presented a higher number of immature, mature, and feeding individuals than the main channel, as well as larger and heavier specimens, mostly caused by the presence of large immature *L. dorsalis* individuals in the creek, and of *A. aspredo* in the main channel, a small sized species. The use of estuarine streams as breeding areas was also observed in Viana *et al.* (2010) and Mourão *et al.* (2015). This difference suggests that the stream environment may offer more suitable conditions for the growth, feeding, and development of juvenile fish (Muller & Strydom 2017, James *et al.* 2019). Preserving and conserving these creeks can be crucial to ensuring successful reproduction and recruitment of new individuals to fish populations.

The freshwater environmental group (FS) was dominant during the study period, followed by estuarine species (ES) that can get around throughout the estuary. This might be explained by the fact that this island presents weak marine influence due to the huge discharge of freshwater, with null salinity but still subject to tidal influence. This riverine dominated system can be classified as a river-mouth estuary (Whitfield & Elliott 2011), where conditions can range from freshwater (<0.5 salinity) to oligohaline waters (0.5 – 4.9 salinity) (Whitfield 2015), which selects freshwater and excludes most estuarine and marine species.

Seven different trophic guilds were observed at Onças Island, wherein zoobenthivores and piscivores predominated throughout the year, corroborating with what is observed at most estuaries (Blaber 2008, Barletta & Blaber 2007, Franco *et al.* 2008, Ferreira *et al.* 2019). Categorizing communities into trophic groups can help simplify the complexity of certain communities and help understand different ecological processes (Austen *et al.* 1994, Benoit *et al.* 2021). In addition, as the number of trophic guilds and their spatial and temporal variations are positively linked to habitat integrity and structure (Elliot *et al.* 2007, Henriques *et al.* 2008, An *et al.* 2013, Mourao *et al.* 2014), the Onças Island can possibly be considered a healthy environment with good integrity.

Plankton carries out an important role in estuaries, which present the bigger productivity in all marine and estuarine ecosystems: the inorganic carbon and nutrients consumed by phytoplankton is linked to the higher trophic levels by zooplankton, a secondary consumer that serves as the major food source for higher trophic species, including commercial fishes (Selleslagh *et al.* 2012, Mitra & Zaman 2016). Accordingly, zoobenthivore species were dominant in all hydrological periods and habitat types in the Onças Island, but the presence of planktivorous species was higher in the tidal creek when compared to the main channel, mostly during dry periods, which is expected, since their main source of food is more abundant at this time of year (Breckenridge *et al.*, 2015). During the dry season, the volume of water inflow into the Guajará bay becomes smaller due to the reduced rainfall, expanding the photic layer and increasing phytoplankton productivity (Lancelot & Muylaert 2011, Cavalcanti *et al.* 2020). Moreover, the highly abundant aquatic

insects, annelids, and crustaceans in Amazonian estuarine environments (da Rocha Leite *et al.* 2013, Filho & Aviz 2013) may use the protection from tidal creeks to spawn (Lucena-Fredou *et al.* 2010, Nóbrega *et al.* 2013; 2014), which may increase the intake of zooplankton by zoobenthivore and planktivore species. Furthermore, planktivore fishes are important to the ecosystem by being the main prey for piscivorous fishes and controlling plankton populations (Soe *et al.* 2021).

Previous studies in other Amazonian estuarine ecosystems have showed changes in the composition of fish communities between seasons (Barthem 1985; Barletta *et al.* 2003; Camargo & Isaac 2001; Hercos 2006, Begot *et al.* 2016), but in Onças Island the variation in the relative abundance of freshwater and estuarine species through the year was small. In conclusion, the Onças Island serves as nursery, reproduction, and feeding area. The results were similar to those found in adjacent areas of Onças Island, such as Guarajá Bay and Guamá River (Mourão *et al.* 2014; Mourão *et al.* 2015; Viana *et al.* 2010). When compared to Amazonian estuaries with greater marine influence, the island showed less diversity because, in these environments, the higher salinity favors a greater diversity of species (Kennish 1986; Marceniuk *et al.* 2021b). The Onças Island is important to the development of several fish species, possibly reflecting the importance of the other islands in Guarajá Bay and other similar Amazon systems, highlighting the need of conservation strategies that reduce anthropogenic impacts on these areas.

ACKNOWLEDGMENTS

We would like to thank the project “Potenciais Impactos Ambientais do Transporte de Petróleo e Derivados na Zona Costeira Amazônica” (PIATAM-Mar) for funding the expeditions in which the samples for this research were collected; the “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior” (CAPES) for the scholarship; and the Instituto Chico Mendes de Conservação da Biodiversidade (ICMbio/SISBio) for the collection license (Authorization #17250-1).

REFERENCES

- Aundsen, P. A., & Sánchez-Hernández, J. (2019). Feeding studies take guts—critical review and recommendations of methods for stomach contents analysis in fish. *Journal of Fish Biology*, 95(6), 1364–1373.
- An, K. G., Choi, J. W., & Lee, Y. J. 2013. Modifications of ecological trophic structures on chemical gradients in lotic ecosystems and their relations to stream ecosystem health. *Animal Cells and Systems*, 17(1), 53–62.
- Austen, D.J., Bayley, P.B., & Menzel, B.W. 1994. Importance of the guild concept to fisheries research and management. *Fisheries*, 19(6): 12–20. doi:10.1577/1548-8446(1994)019<0012:IOTG CT>2.0.CO;2.
- Barletta, M., & Blaber, S. J. M. 2007. Comparison of fish assemblages and guilds in tropical habitats of the Embley (Indo-west Pacific) and caeté (Western Atlantic) estuaries. *Bulletin of Marine Science*, 80(3), 647–680.
- Barletta, M., & Lima, A. R. 2019. Systematic review of fish ecology and anthropogenic impacts in South American estuaries: setting priorities for ecosystem conservation. *Frontiers in Marine Science*, 6, 237.
- Barletta, M., Amaral, C. S., Corrêa, M. F. M., Guebert, F., Dantas, D. V., Lorenzi, L., & Saint-Paul, U. 2008. Factors affecting seasonal variations in demersal fish assemblages at an ecocline in a tropical–subtropical estuary. *Journal of Fish Biology*, 73(6), 1314–1336. DOI: 10.1111/j.1095-8649.2008.02005.x
- Barletta, M., Barletta-Bergan, A., Saint-Paul, U., & Hubold, G. 2003. Seasonal changes in density, biomass, and diversity of estuarine fishes in tidal mangrove creeks of the lower Caeté Estuary (northern Brazilian coast, east Amazon). *Marine Ecology Progress Series*, 256, 217–228. DOI: 10.3354/meps256217
- Barletta, M., Barletta-Bergan, A., Saint-Paul, U., & Hubold, G. 2005. The role of salinity in structuring the fish assemblages in a tropical estuary. *Journal of Fish Biology*, 66(1), 45–72. DOI: 10.1111/j.0022-1112.2005.00582.x
- Barletta, M., Jaureguizar, A. J., Baigun, C., Fontoura, N. F., Agostinho, A. A., Almeida-Val, V. M. F., Val, A. L., Torres, R. A., Jimenes-Segura, L. F., Giarrizzo, T., Fabré, N. N., Batista, V. S., Lasso, C., Taphorn,

- D. C., Costa, M. F., Chaves, P. T., Vieira, J. P., & Corrêa, M. F. M. 2010. Fish and aquatic habitat conservation in South America: a continental overview with emphasis on neotropical systems. *Journal of Fish Biology*, 76(9), 2118–2176. DOI: 10.1111/j.1095-8649.2010.02684.x
- Barros, D. de F., Torres, M. F., & Frédou, F. L. 2011. Ictiofauna do estuário de São Caetano de Odivelas e Vigia (Pará, Estuário Amazônico). *Biota Neotropica*, 11, 367–373. DOI: 10.1590/S1676-06032011000200035
- Barthem, R. B. 1985. Ocorrência, distribuição e biologia dos peixes da Baía de Marajó, estuário amazônico.
- Bartulović, V., Glamuzina, B., Conides, A., Dulčić, J., Lučić, D., Njire, J., & Kožul, V. 2004. Age, growth, mortality and sex ratio of sand smelt, *Atherina boyeri* Risso, 1810 (Pisces: Atherinidae) in the estuary of the Mala Neretva River (middle-eastern Adriatic, Croatia). *Journal of Applied Ichthyology*, 20(5), 427–430.
- Beck, M. W., Heck, K. L., Able, K. W., Childers, D. L., Eggleston, D. B., Gillanders, B. M., Halpern, B. S., Hays, C. G., Hoshino, K., Minello, T. J., Orth, R. J., Sheridan, P. F., & Weinstein, M. P. 2003. The role of nearshore ecosystems as fish and shellfish nurseries. *Issues in Ecology*.
- Beck, Michael W., Heck, K. L., Able, K. W., Childers, D. L., Eggleston, D. B., Gillanders, B. M., Halpern, B., Hays, C. G., Hoshino, K., Minello, T. J., Orth, R. J., Sheridan, P. F., & Weinstein, M. P. 2001. The Identification, Conservation, and Management of Estuarine and Marine Nurseries for Fish and Invertebrates: A better understanding of the habitats that serve as nurseries for marine species and the factors that create site-specific variability in nursery quality will improve conservation and management of these areas. *BioScience*, 51(8), 633–641. DOI:10.1641/0006-3568(2001)051[0633:TICAMO]2.0.CO;2
- Begot, T. O., Soares, B. E., Juen, L., & de Assis Montag, L. F. 2016. Rockpool ichthyofauna of Amazon coastal zone: spatial and environmental effects on species distribution. *Marine and Freshwater Research*, 68(6), 1137–1143.
- Benoit, D. M., Jackson, D. A., & Chu, C. 2021. Partitioning fish communities into guilds for ecological analyses: an overview of current approaches and future directions. *Canadian Journal of Fisheries and Aquatic Sciences*, 78(7), 984–993.
- Bentes, B., Isaac, V. J., Espírito-Santo, R. V. D., Frédou, T., Almeida, M. C. D., Mourão, K. R. M., & Frédou, F. L. 2012. Abordagem multidisciplinar para a identificação dos sistemas de produção pesqueira na costa Norte do Brasil. *Biota Neotropica*, 12, 81–92
- Blaber, S. J. M. 2008. *Tropical Estuarine Fishes: Ecology, Exploitation and Conservation*. John Wiley & Sons: p. 387.
- Blaber, S. J. M. 2013. Fishes and fisheries in tropical estuaries: the last 10 years. *Estuarine, Coastal and Shelf Science*, 135, 57–65.
- Breckenridge, J. K., Bollens, S. M., Rollwagen-Bollens, G., & Roegner, G. C. 2015. Plankton assemblage variability in a river-dominated temperate estuary during late spring (high-flow) and late summer (low-flow) periods. *Estuaries and Coasts*, 38, 93–103.
- Bulger, A. J., Hayden, B. P., Monaco, M. E., Nelson, D. M., & McCormick-Ray, M. G. (1993). Biologically-based estuarine salinity zones derived from a multivariate analysis. *Estuaries*, 16, 311–322.
- Callihan, J. L., Cowan Jr, J. H., & Harbison, M. D. 2013. Sex differences in residency of adult spotted seatrout in a Louisiana estuary. *Marine and Coastal Fisheries*, 5(1), 79–92.
- Camargo, M., & Isaac, V. J. 2001. Os peixes estuarinos da região norte do Brasil: lista de espécies e considerações sobre sua distribuição geográfica.
- Camargo, M., Isaac, V. J., & Fernandes, M. E. B. 2003. Os manguezais da costa norte brasileira. *Fundação Rio Bacanga*, 105–142.
- Castro, P., & Huber, M. E. 2012. *Biologia Marinha* - 8ed. AMGH Editora: p. 480.
- Cavalcanti, L. F., Cutrim, M. V. J., Lourenço, C. B., Sá, A. K. D. S., Oliveira, A. L. L., & de Azevedo-Cutrim, A. C. G. 2020. Patterns of phytoplankton structure in response to environmental gradients in a macrotidal estuary of the Equatorial Margin (Atlantic coast, Brazil). *Estuarine, Coastal and Shelf Science*, 245, 106969. DOI: 10.1016/j.ecss.2020.106969
- Chaudhary, A., Pourfaraj, V., & Mooers, A. O. 2018. Projecting global land use-driven evolutionary history loss. *Diversity and Distributions*, 24(2), 158–167. DOI: 10.1111/ddi.12677
- Chaves, P., Pichler, H., & Robert, M. 2002. Biological, technical and socioeconomic aspects of the fishing activity in a Brazilian estuary. *Journal of Fish Biology*, 61, 52–59.

- Chilton, D., Hamilton, D. P., Nagelkerken, I., Cook, P., Hipsey, M. R., Reid, R., Sheaves, M., Waltham, N. J. & Brookes, J. 2021 Environmental Flow Requirements of Estuaries: Providing Resilience to Current and Future Climate and Direct Anthropogenic Changes. *Front. Environ. Sci.* 9:764218. doi: 10.3389/fenvs.2021.764218
- Costa, M. F., Landing, W. M., Kehrig, H. A., Barletta, M., Holmes, C. D., Barrocas, P. R. G., Evers, D. C., Buck, D. G., Vasconcellos, A. C., Hacon, S. S., Moreira, J. C. & Malm, O. 2012. Mercury in tropical and subtropical coastal environments. *Environmental Research*, 119, 88–100.
- da Rocha Leite, N., Magalhães, A., Palma, M. B., Montes, M. F., Pereira, L. C. C., & da Costa, R. M. 2013. Diel variation in the zooplankton of a highly dynamic Amazonian estuary. *Journal of Coastal Research*, (65), 1146 - 1151.
- Dame, R. F. 2008. Estuaries. *Encyclopedia of Ecology*, 1407-1413. DOI: 10.1016/b978-008045405-4.00329-3
- Dantas, D. V., Barletta, M., Costa, M. F., Barbosa-Cintra, S. C. T., Possatto, F. E., Ramos, J. A., ... & Saint-Paul, U. (2010). Movement patterns of catfishes (Ariidae) in a tropical semi-arid estuary. *Journal of Fish Biology*, 76(10), 2540-2557.
- De Araujo-Barbosa, C. C., Atkinson, P. M., & Dearing, J. A. 2016. Extravagance in the commons: Resource exploitation and the frontiers of ecosystem service depletion in the Amazon estuary. *Science of The Total Environment*, 550, 6–16. DOI: 10.1016/j.scitotenv.2016.01.072
- Dias, J. A., Cearreta, A., Isla, F. I., & de Mahiques, M. M. 2013. Anthropogenic impacts on Iberoamerican coastal areas: Historical processes, present challenges, and consequences for coastal zone management. *Ocean & Coastal Management*, 77, 80–88. DOI: 10.1016/j.ocecoaman.2012.07.025
- Dunn, O. J. 1961. Multiple comparisons among means. *JASA*, 56: 54 - 64.
- Earl, J., Fowler, A. J., Ye, Q., & Dittmann, S. 2014. Age validation, growth and population characteristics of greenback flounder (*Rhombosolea tapirina*) in a large temperate estuary. *New Zealand Journal of Marine and Freshwater Research*, 48(2), 229-244.
- Elliott, M., Whitfield, A. K., Potter, I. C., Blaber, S. J. M., Cyrus, D. P., Nordlie, F. G., & Harrison, T. D. 2007. The guild approach to categorizing estuarine fish assemblages: a global review. *Fish and Fisheries*, 8(3), 241–268. DOI: 10.1111/j.1467-2679.2007.00253.x.
- Ferreira, V., Le Loc'h, F., Ménard, F., Frédou, T., & Frédou, F. 2019. Composition of the fish fauna in a tropical estuary: the ecological guild approach. *Scientia Marina*, 83(2), 133–142. DOI: 10.3989/scimar.04855.
- Filho, J. S. R., & Aviz, D. Macrobenthic communities of an Amazonian estuary (Guajará Bay, Brazil): temporal and spatial changes. 2013. *Journal of Coastal Research*, (65), 123-128.
- Firth, L., Knights, A., Bridger, D., Evans, A., Mieszkowska, N., Moore, P., O'Connor, N., Sheehan, E., Thompson, R., & Hawkins, S. 2016. Ocean sprawl: challenges and opportunities for biodiversity management in a changing world. *Oceanography and Marine Biology*, 54, 189–262.
- Franco, A, Elliott, M., Franzoi, P., & Torricelli, P. 2008. Life strategies of fishes in European estuaries: the functional guild approach. *Marine Ecology Progress Series*, 354(2), 219–228. DOI: 10.3354/meps07203
- Froese, R., & Pauly, D. 2018. Search FishBase. (Retrieved on October 29th, 2023, from <https://www.fishbase.se/search.php>).
- Gregório, A. M. S., & Mendes, A. C. 2009. Batimetria e sedimentologia da baía de guajará, belém, estado do pará, brasil. *Amazônia: Ci. & Desenv.*, 5(9), 53–72.
- Guebert-Bartholo, F. M., Barletta, M., Costa, M. F., Lucena, L. R., & da Silva, C. P. 2011. Fishery and the use of space in a tropical semi-arid estuarine region of Northeast Brazil: subsistence and overexploitation. *Journal of Coastal Research*, 398-402.
- Harrison, T., & Whitfield, A. 2006. Temperature and salinity as primary determinants influencing the biogeography of fishes in South African estuaries. *Estuarine, Coastal and Shelf Science*, 66, 335–345. DOI: 10.1016/j.ecss.2005.09.010
- Hashiguti, D. T., Rocha, R. M., & Montag, L. F. Reproductive seasonality of the detritivorous fish *Cyphocharax abramoides* (Kner, 1958) (Characiformes: Curimatidae) in flooded rivers of the eastern Amazon. 2017. *Environmental Biology of Fishes*, 100, 1033-1046.
- Hashiguti, D. T., Soares, B. E., Wilson, K. L., Oliveira-Raiol, R. D., & Montag, L. F. D. A. Comparing three methods to estimate the average size at first maturity: a case study on a Curimatid exhibiting polyphasic growth. 2019. *Ecology of Freshwater Fish*, 28(2), 266-273.

- Healy, T. R. 2005. Tidal Creeks. In: Schwartz, M.L. (eds) Encyclopedia of Coastal Science. Encyclopedia of Earth Science Series. Springer, Dordrecht. https://doi.org/10.1007/1-4020-3880-1_313
- Henriques, S., Pais, M. P., Costa, M. J., & Cabral, H. 2008. Development of a fish-based multimetric index to assess the ecological quality of marine habitats: the Marine Fish Community Index. *Marine Pollution Bulletin*, 56(11), 1913–1934. DOI: 10.1016/j.marpolbul.2008.07.009
- Hercos, A. P. 2006. Diversidade e variabilidade espaço-temporal da ictiofauna da região estuarina do rio Curuçá município de Curuçá, Pará Brasil.
- Hill, M. O. 1973. Diversity and evenness: a unifying notation and its consequences. *Ecology*, 54(2), 427-432.
- IBGE. 2022. Instituto Brasileiro de Geografia e Estatística - IBGE. 2022. (Retrieved on November 5th, 2023, from <https://cidades.ibge.gov.br/brasil/pa/belem/panorama>).
- INMET. 2013. (Retrieved on June 1st, 2006, from <http://www.inmet.gov.br/>). Instituto Nacional de Meteorologia - INMET. 2013. (Retrieved on October 28th, 2023, from <https://portal.inmet.gov.br/noticias/noticias?noticias=2013>).
- James, N. C., Leslie, T. D., Potts, W. M., Whitfield, A. K., & Rajkaran, A. 2019. The importance of different juvenile habitats as nursery areas for a ubiquitous estuarine-dependent marine fish species. *Estuarine, Coastal and Shelf Science*, 226, 106270. DOI: 10.1016/j.ecss.2019.106270
- Kennish, M. 1986. *Ecology of Estuaries, Volume 1 Physical and Chemical Aspects*. p. book.
- Kennish, M. J. 2002. Environmental threats and environmental future of estuaries. *Environmental conservation*, 29(1), 78-107
- Kolpakov, N. V., & Milovankin, P. G. 2010. Distribution and seasonal changes in fish abundance in the estuary of the Razdol'naya River (Peter the Great Bay), Sea of Japan. *Journal of Ichthyology*, 50(6), 445-459.
- Lancelot, C., & Muylaert, K. 2011. Trends in Estuarine Phytoplankton Ecology. *Treatise Estuar Coast Sci*, 7, 5–15. DOI: 10.1016/B978-0-12-374711-2.00703-8
- Latrubesse, E., Arima, E., Dunne, T. *et al.* 2017. Damming the rivers of the Amazon basin. *Nature* 546, 363–369. DOI: 10.1038/nature22333
- Lessa, G. C., Santos, F. M., Souza Filho, P. W., & Corrêa-Gomes, L. C. 2018. Brazilian estuaries: a geomorphologic and oceanographic perspective. *Brazilian Estuaries: A Benthic Perspective*, 1-37.
- Lucena-Fredou, F., Rosa Filho, J. S., Silva, M. C., & Azevedo, E. F. 2010. Population dynamics of the river prawn, *Macrobrachium amazonicum* (Heller, 1862)(Decapoda, Palaemonidae) on Combu Island (Amazon estuary). *Crustaceana*, 83(3): 277-290.
- Marceniuk, A. P., Caires, R. A., Carvalho-Filho, A., de Macedo Klautau, A. G. C., Santos, W. C. R., Wosiacki, W. B., ... & Rotundo, M. M. 2021a. Teleostei fishes of the North Coast of Brazil. *Revista CEPISUL-Biodiversidade e Conservação Marinha*, 10, e2021006-e2021006
- Marceniuk, A. P., Caires, R. A., Carvalho-Filho, A., Rotundo, M. M., Santos, W. C. R. D., & Klautau, A. G. C. D. M. 2021b. Peixes teleósteos da costa norte do Brasil. *Museu Paraense Emílio Goeldi*.
- Marceniuk, A. P., Soares, B. E., Rotundo, M. M., Caires, R. A., Rosa, R. D. S., Santos, W. C. R. D. & Barthém, R. 2023. The bycatch of piramutaba, *Brachyplatystoma vaillantii* industrial fishing in a salinity and depth gradient in the Amazon estuary, Brazil. *Acta Amazonica*, 53, 93-106.
- McLusky, D. S., & Elliot, M. The estuarine ecology: ecology, threats and management. 2004. Oxford University Press, UK.
- Merigot, B., Frédou, F. L., Viana, A. P., Ferreira, B. P., Junior, E. D. N. C., da Silva Júnior, C. B., & Frédou, T. 2017. Fish assemblages in tropical estuaries of northeast Brazil: A multi-component diversity approach. *Ocean & coastal management*, 143, 175-183.
- Mitra, A., & Zaman, S. (2016). *Basics of marine and estuarine ecology*. Springer.
- Molina, A., Duque, G., & Cogua, P. 2020. Influences of environmental conditions in the fish assemblage structure of a tropical estuary. *Marine Biodiversity*, 50(1), 5.
- Mourão, K. R. M., Frédou, T., & Frédou, F. L. 2015. Spatial and seasonal variation of the ichthyofauna and habitat use in the inner portion of the Brazilian Amazon estuary. *Boletim Do Instituto de Pesca*, 41(3), 529–545.
- Mourão, K. R., Ferreira, V., & Lucena-Fredou, F. 2014. Composition of functional ecological guilds of the fish fauna of the internal sector of the Amazon Estuary, Pará, Brazil. *Anais da Academia Brasileira de Ciências*, 86, 1783-1800.

- Muller, C., & Strydom, N. A. 2017. Evidence for Habitat Residency and Isotopic Niche Partitioning in a Marine-Estuarine-Dependent Species Associated with Mangrove Habitats from the East Coast of South Africa. *Estuaries and Coasts*, 40(6), 1642–1652. DOI: 10.1007/s12237-017-0240-3
- Nóbrega, P. S. V. D., Bentes, B., & Martinelli-Lemos, J. M. Composition of shrimp populations (Crustacea: Decapoda) in non-vegetated areas of two river islands in a Brazilian Amazon estuary. 2013. *Zoologia (Curitiba)*, 30, 652-660.
- Nóbrega, P. S. V. D., Bentes, B., & Martinelli-Lemos, J. M. Population structure and relative growth of the Amazon shrimp *Macrobrachium amazonicum* (Heller, 1862)(Decapoda: Palaemonidae) on two islands in the fluvial-estuarine plain of the Brazilian Amazon. 2014. *Nauplius*, 22, 13-20.
- Nunes, T. G., & Furtado, L. de F. G. 2023. A ilha do Combu: ensaio sobre turismo e lazer em intenso crescimento. Version 1. *Novos Cadernos NAEA*, 26(1). DOI: 10.18542/ncn.v26i1.11121
- Núñez, J., & Duponchelle, F. Towards a universal scale to assess sexual maturation and related life history traits in oviparous teleost fishes. 2009. *Fish physiology and biochemistry*, 35, 167-180.
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. *vegan: Community ecology package*. 2019. R package version 2.5-4.
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. 2007. Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, 11(5), 1633–1644. DOI: 10.5194/hess-11-1633-2007
- Pichler, H. A., Spach, H. L., Gray, C. A., Broadhurst, M. K., Schwarz Jr, R., & de Oliveira Neto, J. F. 2015. Environmental influences on resident and transient fishes across shallow estuarine beaches and tidal flats in a Brazilian World Heritage area. *Estuarine, Coastal and Shelf Science*, 164, 482-492.
- Pinheiro, F. G. R. 2002. Avaliação dos riscos de contaminação dos aquíferos livres da bacia do Tucunduba /Belém (PA). 2002. 55 f. Trabalho de Conclusão de Curso (Graduação em Geologia) – Centro de Geociências, Universidade Federal do Pará. Belém. Porto de Vila do Conde – Portal CDP. 2023.
- Potter, I. C., Chuwen, B. M., Hoeksema, S. D., Elliott, M. 2010. The concept of an estuary: A definition that incorporates systems which can become closed to the ocean and hypersaline. *Estuarine, Coastal and Shelf Science*. 87, 497-500.
- Prestes, L., Barthem, R., Mello-Filho, A., Anderson, E., Correa, S. B., Couto, T. B. D. A., ... & Goulding, M. Proactively averting the collapse of Amazon fisheries based on three migratory flagship species. 2022. *Plos one*, 17(3), e0264490.
- R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing.
- Raz-Guzman, A. & Huidobro, L. 2002. Fish communities in two environmental different estuarine systems of Mexico. *J. Peixe Biol.* 60, 1–14.
- Ribeiro, K. T. S. 2004. Água e saúde humana em Belém. Belém: Cejup, 2004. 280 p. (Coleção Megam).
- Rundle, S. D., M. J. Attrill & A. Arshad. 1998. Seasonality in macroinvertebrate community composition across a neglected ecological boundary, the freshwater–estuarine transition zone. *Aquatic Ecology* 32, 211–216.
- Secor, D. H. 1999. Specifying divergent migrations in the concept of stock: the contingent hypothesis. *Fisheries Research*, 43(1-3), 13-34.
- Selleslagh, J., Amara, R., Laffargue, P., Lesourd, S., Lepage, M., & Girardin, M. 2009. Fish composition and assemblage structure in three Eastern English Channel macrotidal estuaries: A comparison with other French estuaries. *Estuarine, Coastal and Shelf Science*, 81(2), 149–159. DOI: 10.1016/j.ecss.2008.10.008
- Selleslagh, J., Lobry, J., Amara, R., Brylinski, J. M., & Boët, P. 2012. Trophic functioning of coastal ecosystems along an anthropogenic pressure gradient: a French case study with emphasis on a small and low impacted estuary. *Estuarine, Coastal and Shelf Science*, 112, 73-85.
- Silva, A. C. B., Barros, M. S. F., Silva, V. E. L., de Oliveira, C. D. L., Santos, M. E. F., & Fabr , N. N. 2023. Composition and distribution of fish assemblages in a tropical river–estuarine continuum. *Hydrobiologia*, 1-11.
- Soares, B. E., Barros, T. F., Hashiguti, D. T., Pereira, D. C., Ferreira, K. C., & Caramaschi,  . P. 2020. Traditional approaches to estimate length at first maturity (L 50) retrieve better results than alternative ones in a Neotropical heptapterid. *Journal of Fish Biology*, 97(5), 1393-1400.

- Soares, B. E., Benone, N. L., Barthem, R. B., Marceniuk, A. P., & Montag, L. F. 2021. Environmental conditions promote local segregation, but functional distinctiveness allows aggregation of catfishes in the Amazonian estuary. *Estuarine, Coastal and Shelf Science*, 251, 107256.
- Soe, K. K., Pradit, S., & Hajisamae, S. 2021. Feeding habits and seasonal trophic guilds structuring fish community in the bay mouth region of a tropical estuarine habitat. *Journal of Fish Biology*, 99(4), 1430-1445.
- Van der Sleen, P., & Albert, J. S. (Eds.). (2017). *Field guide to the fishes of the Amazon, Orinoco, and Guianas* (Vol. 115). Princeton University Press.
- Vergès, L. H. M. C., Contente, R. F., Marion, C., Castillo, C. P. C. D., Spach, H. L., Cattani, A. P., & Fávaro, L. F. (2022). Relationship between fish assemblage structure and predictors related to estuarine productivity in shallow habitats of a Neotropical estuary. *Neotropical Ichthyology*, 20.
- Viana, A. P., & Lucena Frédou, F. 2014. Ichthyofauna as bioindicator of environmental quality in an industrial district in the amazon estuary, Brazil. *Brazilian Journal of Biology*, 74, 315–324. DOI: 10.1590/1519-6984.16012.
- Viana, A. P., Lucena Frédou, F., Frédou, T., Torres, M. F., & Bordalo, A. O. 2010. Fish fauna as an indicator of environmental quality in an urbanised region of the Amazon estuary. *Journal of Fish Biology*, 76(3), 467–486. DOI: 10.1111/j.1095-8649.2009.02487.x.
- Walsh, C. T., Pease, B. C., & Booth, D. J. 2004. Variation in the sex ratio, size and age of longfinned eels within and among coastal catchments of south-eastern Australia. *Journal of Fish Biology*, 64(5), 1297-1312.
- Whitfield, A. K. 2015. Why are there so few freshwater fish species in most estuaries?. *Journal of Fish Biology*, 86(4), 1227-1250.
- Whitfield, A. K., & Harrison, T. D. 2003. River flow and fish abundance in a South African estuary. *Journal of Fish Biology*, 62(6), 1467-1472.
- Whitfield, A. K., Elliott, M., Basset, A., Blaber, S. J. M., & West, R. J. 2012. Paradigms in estuarine ecology—a review of the Remane diagram with a suggested revised model for estuaries. *Estuarine, Coastal and Shelf Science*, 97, 78-90.
- Whitfield, A.K., & Elliot, M. 2011. Ecosystem and biotic classifications of estuaries and coasts. In: Wolanski, E. & McLusky, D. (Eds), *Treatise on Estuarine and Coastal Science*. pp. 99-124. Academic Press.
- Zar, J. D H. 1996. *Biostatistical analysis*. 3 edição. Ed. Prentice-Hall International INC. 662 p.

SUPPLEMENTARY MATERIAL

Appendix A. Monthly general data of precipitation, richness, abundance, and evenness of fish collected between September 2006 and September 2008, Onças Island, Pará, Brazil.

Appendix B. Rarefaction analysis presenting the interpolation with observed data (solid lines) and extrapolation (dotted lines) curves using Hill-number for species richness ($q = 0$) by habitat type (Main Channel, Tidal Creek, and the Onças Island), Pará, Brazil. Shaded areas represent 95 % confidence intervals.

Appendix C. Total length frequency of captured fish, presented by class intervals (5cm), between September 2006 and September 2008, Onças Island, Pará, Brazil.

Appendix D. Percentage of individuals with any amount of visible organic material in the stomach (With food) for each feeding guild (CV: Carnivore, OV: Omnivore, PV: Piscivore, HV: Herbivore, ZB: Zoobenthivore, PL: Planktivore; DV: Detritivore), in fish collected between September 2006 and September 2008, Onças Island, Pará, Brazil.

Submitted: 27 September 2023

Accepted: 04 May 2024

Published online: 16 May 2024

Associate Editor: Bruno Soares