



RECORD OF VISIBLE IMPLANT ALPHA TAG LOSS IN THE CATFISH *Rhamdioglanis transfasciatus* (SILURIFORMES: HEPTATERIDAE) INHABITING AN ATLANTIC FOREST STREAM

Thiago Fonseca de Barros¹ , Bruno Eleres Soares^{2*} ,
Marcus Vinícius Vieira³ & Érica Pellegrini Caramaschi⁴

¹ Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Ecologia, Programa de Pós-Graduação em Ecologia, Laboratório de Ecologia de Peixes, Av. Carlos Chagas Filho, 373, 21941-902, Rio de Janeiro, Rio de Janeiro, Brazil.

² University of Regina, Faculty of Science, Institute of Environmental Change & Society, Department of Biology, Soares Quantitative Ecology Lab, 3737 Wascana Parkway, S4S0A2, Regina, Saskatchewan, Canada.

³ Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Ecologia, Programa de Pós-Graduação em Ecologia, Laboratório de Vertebrados, Av. Carlos Chagas Filho, 373, 21941-902, Rio de Janeiro, Rio de Janeiro, Brazil.

⁴ Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Ecologia, Programa de Pós-Graduação em Ecologia, Laboratório de Ecologia de Peixes, Av. Carlos Chagas Filho, 373, 21941-902, Rio de Janeiro, Rio de Janeiro, Brazil.

E-mails: tod_barros@yahoo.com.br; bruno.soares@uregina.ca (*corresponding author); mvviera@gmail.com; erica.caramaschi@gmail.com

Abstract: We report the number of recaptured individuals and the loss of visible implant alpha tags in the Neotropical catfish *Rhamdioglanis transfasciatus* during a mark-recapture study conducted in Atlantic Forest streams, Brazil. We recaptured 1.8% and 29% of the marked individuals in the rainy and dry seasons, respectively. The lowest number of recaptured labels occurred in the rainy, warm season, which could be attributed both to higher movement events of individuals outside the monitored area or because the inflammatory reaction that leads to tag loss, which is favored in warmer conditions. These findings highlight the need to improve the application of VI Alpha tags, whether through refining tagging protocols and materials, so as to enhance tracking accuracy and minimize resource loss in ecological studies.

Keywords: Atlantic Forest; freshwater fish; mark-capture; stream; tagging.

Marking fish individuals is fundamental to understanding ecological patterns related to movement, growth, survival, and habitat use. Common techniques include fin clipping, chemical or dye marking, external tags, passive integrated transponders, and elastomer-based markers. Those methods are associated to different levels of physical injuries (e.g. Ivasauskas et al. 2012, Collins et al. 2013),

fitness reduction (e.g. Gauthier-Clerc et al. 2004, Wilson et al. 2011; Ouedraogo et al. 2014), predation risks (e.g. Carlson & Langkilde 2013), and mortality rates (e.g. Økland et al. 2001, Gauthier-Clerc et al. 2004, Wilson et al. 2011). In stream environments, where fish are often small-bodied and habitats are structurally complex, researchers frequently use visible implant elastomer (VIE) tags and visible

implant alpha (VI Alpha) tags to individually mark fish (e.g., Mazzoni & Iglesias-Rios 2012, Mazzoni et al. 2018). VIE tags pose issues for individualize members of the population, thus hampering the obtention of individual-level information on growth and condition (Jungwirth et al. 2019). In turn, externally VI Alpha tags are small fluorescent tags inscribed with unique alphanumeric codes (Knapp et al. 2023), making them suitable for tagging small-sized fishes given their size, easiness of application, and low cost (Turek et al. 2014).

In addition to the negative effects to the organism (e.g., mortality, injuries), there are many records of expulsion of internally implanted markers, such as subcutaneous labels and passive integrated transponders (Jepsen et al. 2002). Studies assessing the tag retention rates of VI alpha tags (Northwest Marine Technology, Inc.) have shown a wide range of variation in retention (Turek et al. 2014). Tag expulsion does not compromise the organism fitness and survival (Jepsen et al. 2002), but poses a loss of time, material, resources and information to

the population study. Herein, we report the loss of VI alpha tags in the Neotropical catfish *Rhamdioglanis transfasciatus*, a heptapterid inhabiting Atlantic Forest streams, following a mark-recapture experiment. We carried out four expeditions in the rainy season and four expeditions in the dry season of 2016, each consecutive expedition within the season in a 20-day interval.

We captured fish by electrofishing (Mazzoni et al. 2000) within a 375 m stretch in the “Mineiro-Branco” Stream (Figure 1; 22° 25' 18.26" S, 42° 44' 20.90" W), Guapiaçu River basin, within a preserved Atlantic Forest area in the Rio de Janeiro State, southeastern Brazil. We used an alternate current generator (Honda EU10i, 1.000W, 220V, 4.1A), with two electrified poles. After stunning, individuals were captured from the water column using nonelectrified hand nets. Captured individuals were kept in floating cages, with water circulation from the stream itself, for total recovering from electronarcosis. More details on sampling techniques can be found in Soares et al. (2020).




Figure 1. Location of the Mineiro-Branco Stream, located in the Guapiaçu River basin, southeastern Brazil.

Sampled individuals were anesthetized with Eugenol and marked through an incision in the ventral region of the caudal peduncle with VI Alpha Tags, biocompatible labels individualized by color and alphanumeric code. More details are available in the manufacturer's website (<http://www.nmt.us/vi-alpha/>). We used a total of 842 labels (Table 1) and did not record any mortality due to anesthesia, handling and/or marking procedures.

During the field expeditions, we noticed that many fish have lost their labels by the presence of a round-whitish scar in the incision point. Some fish presented advanced inflammatory reactions in the incision location and still conserved their tags (Figure 2). We counted all the times we observed the inflammation process in the incision point for the following descriptive statistics (Table 2): the total number of used labels, the number of

Table 1. Number of individuals caught at each mark-recapture event in the Rainy and Dry seasons of 2016 in the Mineiro-Branco Stream, Rio de Janeiro State, southeastern Brazil.

Event	Rainy/2016	Dry/2016
Day 0	109	104
Day 20	94	89
Day 40	136	101
Day 60	114	95

individuals recaptured with labels, the number of never-recovered labels (= the total number of used labels minus the recovered ones). Proportions of these numbers were calculated as percentages from the total number of used labels. We also calculated the estimated number of labels used per individual within a season by dividing the total number of



Figure 2. Perceptible inflammatory reaction as a purple-colored skin lesion (circle) around the label in an individual of *Rhamdioglanis transfasciatus* collected in the Mineiro-Branco Stream, Rio de Janeiro State, southeastern Brazil.

Table 2. Numbers of recovered and not recovered labels, proportions based on the total number of labels used and the average number of labels used per individual in each season in the Mineiro-Branco Stream, Rio de Janeiro State, southeastern Brazil. The second and last sampling expedition occurred in the dry season, so there is no record of individuals recaptured in its next season.

Season	Rainy		Dry	
	N	% from total	N	% from total
Individuals recaptured with labels	8	1.8%	83	29.0%
Individuals recaptured with labels in the next season	3	0.7%	-	-
Never-recovered labels	431	97.5%	203	71.0%
Maximum number of individuals marked in a season	136	31.6%	104	51.2%
Used labels (Total)	442	100.0%	286	100.0%
Average number of labels used per individual	3.01		1.53	

labels used by the sum of individuals recaptured with labels. One should note that those calculations consider that all individuals displaying labels were marked and recaptured just once, which potentially introduce biases in this description, since some individuals were probably marked more than once.

During the rainy season (i.e., summer in Brazil), we deployed approximately 442 labels, but only 11 tagged individuals were recaptured, representing a recapture rate of 2.5%. Given the low recapture rate, we estimate that more than three labels were used, on average, for each individual fish captured in this season. As the rainy season corresponded to the first sampling expedition, the low recapture rate raised concerns about the feasibility and validity of continuing this study. We estimate that about US\$500 was lost in material during this season alone. Differently, the dry season (i.e., winter in Brazil) was more promising in relation to the maintenance of labels in individuals, since 29.0% of all used labels were indeed recovered. The tag losses posed some methodological limitations to the application of the data to understand population dynamics and movement ecology. First, we were unable to determine the exact number of marked individuals during the study. Some individuals may have lost their tags and, without visible signs of tagging or inflammation, were indistinguishable from unmarked individuals. Second, due to the tag loss, we could not accurately calculate population sizes and other population-level descriptors, as necessary parameters (total number of recaptured

individuals at each sampling event) were compromised by low tag retention rates.

Cases of implanted markers' expulsion have already been recorded for other species, including other catfishes such as *Ictalurus punctatus* (Marty & Summerfelt 1986) in the United States and *Rhamdia quelen* (Schulz 2003) in Brazil. Encapsulation prior to expulsion occurs due to the organism's inflammatory reaction to a foreign body, and its subsequent expulsion usually occurs via the incision point or through the body wall (Coleman et al. 1974, Moore et al. 1990, Jepsen et al. 2002). The risk of expulsion is usually reduced if the marker is cylindrical or has rounded tips but increased if it has sharp edges (Thorstad et al. 2013). VI Alpha Tags have a rectangular shape with rounded edges, but are flat and slightly hardened, which gives them relatively sharp edges, thus increasing the risk of expulsion. In addition, we also consider that the lower temperatures during the winter potentially decreased the incidence of inflammatory processes and subsequent tag expulsion. The water temperature ranged from 22.2 to 22.8 °C (average = 22.5 °C, SD = 0.29) in the rainy season and from 16.3 to 18.7 °C (average = 17.8 °C, SD = 1.06) in the dry season. It is reported in the literature that greater infection risk may be linked to the release of fish in relatively warmer waters (Jepsen et al. 2002). Also warmer temperatures promoted tag expulsion in the bluegill sunfish *Lepomis macrochirus* (Knights & Lasee 1996), the striped bass *Morone saxatilis* × white bass *M. chrysops* hybrid (Walsh et al. 2000),

and the chinook salmon *Oncorhynchus tshawytscha* (Deters et al. 2010) in North America.

Studies regarding tag expulsion in Neotropical fish have been conducted under manipulative conditions (e.g. Schulz 2003, Lopes et al. 2016). Although recent studies using biocompatible markers in Neotropical stream fishes (e.g. Mazzoni & Iglesias-Rios 2012, Espírito-Santo et al. 2016, Celestino et al. 2017, Mazzoni et al. 2018) have not reported marker expulsion, several of them were conducted under similar environmental conditions to our study. Therefore, we do not attribute the high tag loss in our study to abiotic factors alone. Instead, we suggest potential explanations: (1) inflammation or scarring may have gone unnoticed by researchers, and (2) interspecific differences in tag retention. Tagging methods exhibit varying levels of tag loss, but they commonly present some losses. Passive integrated transponder tags have shown to present low retention rate in body cavity (Mamer & Meyer 2016) while visible implant elastomer exhibits higher retention rates (Hohn & Petrie-Hanson 2013, Bushon & Rash 2021). While the latter exhibits higher retention rates than the method we used, there are limitations on the number of individuals that can receive a unique combination of colors that allow the recognition of different individuals.

One way to avoid possible premature study interruptions and hindrances caused by loss of time, material, resources, and marked individuals is to use alternative methods of individual identification. Methods such as the detection of electric fields (e.g., Henninger et al. 2020 with Gymnotiformes) or the use of natural skin marks (e.g. Dala-Corte et al. 2016 with Loricariidae) can be easily applied because of currently available technological resources. Despite this, these methods are taxon-dependent and rarely applied, requiring more investigation. Other suggestions would be combining both VI Alpha tags and VIE for better controlling the number of marked individuals and carrying out a pilot tagging program in a more controlled environment to assess retention rates in the focal species. Finally, despite the high loss of labels, the number of marked individuals allowed us to continue the study and promoted novel findings regarding the movement of Neotropical stream-dwelling fish.

ACKNOWLEDGEMENTS

We are grateful to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001 for the master scholarship provided to TFB during the execution of this study. We are also thankful to colleagues in Laboratório de Ecologia de Peixes (UFRJ) for helping in the field expeditions, and to Nicholas J. Locke, Raquel P. Locke, Jorge Bizarro and all employees from the Reserva Ecológica de Guapiaçu (REGUA).

REFERENCES

- Bushon, A. M., & Rash, J. M. 2021. Retention of postocular visible implant elastomer in two sizes of adult brown trout and rainbow trout. *North American Journal of Fisheries Management*, 41(5), 1384-1389. DOI: 10.1002/nafm.10656
- Carlson, B. E., & Langkilde, T. 2013. A common marking technique affects tadpole behavior and risk of predation. *Ethology*, 119(2), 167-177. DOI: 10.1111/eth.12050
- Celestino, L. F., Sanz-Ronda, F. J., Kashiwaqui, E. A. L., Celestino, E. F., Makrakis, M. C., & Makrakis, S. 2017. Daily movement behavior of two Neotropical armored catfish species (*Ancistrus* aff. *cirrhosus* Valenciennes, 1836 and *Hypostomus ancistroides* Ihering, 1911) at a road-stream crossing culvert. *Journal of Applied Ichthyology*, 33(6), 1092-1099. DOI: 10.1111/jai.13446
- Coleman, D. L., King, R. N., & Andrade, J. D. 1974. The foreign body reaction: a chronic inflammatory response. *Journal of Biomedical Materials Research*, 8(5), 199-211. DOI: 10.1002/jbm.820080503
- Collins, A. L., Hinch, S. G., Welch, D. W., Cooke, S. J., & Clark, T. D. 2013. Intracoelomic acoustic tagging of juvenile sockeye salmon: swimming performance, survival, and postsurgical wound healing in freshwater and during a transition to seawater. *Transactions of the American Fisheries Society*, 142(2), 515-523. DOI: 10.1080/00028487.2012.743928
- Dala-Corte, R. B., Moschetta, J. B., & Becker, F. G. 2016. Photo-identification as a technique for recognition of individual fish: a test with the freshwater armored catfish *Rineloricaria*

- aequalicuspis* Reis & Cardoso, 2001 (Siluriformes: Loricariidae). Neotropical Ichthyology, 14(1): e150074. DOI: 10.1590/1982-0224-20150074
- Deters, K. A., Brown, R. S., Carter, K. M., & Boyd, J. W. 2010. Performance assessment of suture type, water temperature, and surgeon skill in juvenile chinook salmon surgically implanted with acoustic transmitters. Transactions of the American Fisheries Society, 139(3), 888-899. DOI: 10.1577/T09-043.1
- Espírito-Santo, H. M. V., Rodríguez, M. A., & Zuanon, J. 2016. Strategies to avoid the trap: stream fish use fine-scale hydrological cues to move between the stream channel and temporary pools. Hydrobiologia, 792(1), 183-194. DOI: 10.1007/s10750-016-3054-6
- Gauthier-Clerc, M., Gendner, J.-P., Ribic, C. A., Fraser, W. R., Woehler, E. J., Descamps, S., Gilly, C., Le Bohec, C., & Le Maho, Y. 2004. Long-term effects of flipper bands on penguins. Proceedings of the Royal Society London B, 271(Suppl. 6), S423-S426. DOI: 10.1098/rsbl.2004.0201
- Henninger, J., Krahe, R., Sinz, F., & Benda, J. 2020. Tracking activity patterns of a multispecies community of gymnotiform weakly electric fish in their neotropical habitat without tagging. Journal of Experimental Biology, 223(3), jeb206342. DOI: 10.1242/jeb.206342
- Hohn, C., & Petrie-Hanson, L. 2013. Evaluation of visible implant elastomer tags in zebrafish (*Danio rerio*). Biology Open, 2(12), 1397-1401. DOI: 10.1242/bio.20136460
- Ivasauskas, T. J., Bettoli, P. W., Holt, T. 2012. Effects of suture material and ultrasonic transmitter size on survival, growth, wound healing, and tag expulsion in rainbow trout. Transactions of the American Fisheries Society, 141(1), 100-106. DOI: 10.1080/00028487.2011.651553
- Jepsen, N., Koed, A., Throstad, E. B., & Baras E. 2002. Surgical implantation of telemetry transmitters in fish: how much have we learned? Hydrobiologia, 483(1), 239-248. DOI: 10.1023/A:1021356302311
- Jungwirth, A., Balzarini, V., Zöttl, M., Salzmann, A., Taborsky, M., & Frommen, J. G. 2019. Long-term individual marking of small freshwater fish: the utility of Visual Implant Elastomer tags. Behavioural Ecology and Sociobiology, 73(1), 49. DOI: 10.1007/s00265-019-2659-y
- Knapp, D. D., Diaz, L., Unger, S., Anderson, C. N., Spear, S. F., Williams, L. A., Kidd-Weaver, A. D., Green, M. W., Poelmann, O. M., Kerr, J., & Bodinof, C. M. 2023. Long-term retention, readability, and health effects of Visible Implant Elastomer (VIE) and Visible Implant Alpha (VI Alpha) tags in larval Eastern Hellbenders (*Cryptobranchus alleganiensis alleganiensis*). Journal of Herpetology, 57(2), 133-141. DOI: 10.1670/22-011
- Knights, B. C., & Lasee B. A. 1996. Effects of implanted transmitters on adult bluegills at two temperatures. Transactions of the American Fisheries Society, 125(3): 440-449. DOI:10.1577/1548-8659(1996)125<0440:EOITO A>2.3.CO;2
- Lopes, J. M., Alves, C. B. M., Silva, F. O., Bedore, A. G., & Pompeu, P. S. 2016. Effect of anesthetic, tag size, and surgeon experience on postsurgical recovering after implantation of electronic tags in a neotropical fish: *Prochilodus lineatus* (Valenciennes, 1837) (Characiformes: Prochilodontidae). Neotropical Ichthyology, 14(3), e150189. DOI: 10.1590/1982-0224-20150189
- Mamer, E. R. J. M., & Meyer, K. A. 2016. Retention rates of passive integrated transponder tags, visible implant elastomer tags, and maxillary marks in wild trout. North American Journal of Fisheries Management, 36(5), 1119-1124. DOI: 10.1080/02755947.2016.1198288
- Marty, G. D., & Summerfelt, R. C. 1986. Pathways and mechanisms for expulsion of surgically implanted dummy transmitters from channel catfish. Transactions of the American Fisheries Society, 115(4), 577-589. DOI:10.1577/1548-8659(1986)115<577:PAMFEO>2.0.CO;2
- Mazzoni, R., Fenerich-Verani, N., & Caramaschi, E. P. 2000. Electrofishing as a sampling technique for coastal stream fish populations and communities in the Southeast of Brazil. Revista Brasileira de Biologia, 60(2), 205-206. DOI: 10.1590/S0034-71082000000200003
- Mazzoni, R., & Iglesias-Rios, R. 2012. Movement patterns of stream-dwelling fishes from Mata Atlântica, Southeast Brazil. Revista de Biologia Tropical, 60(4), 1837-1846.
- Mazzoni, R., Pinto, M. P., Iglesias-Rios, R., & Costa, R. 2018. Fish movement in an Atlantic Forest stream. Neotropical Ichthyology, 16(1), e170065. DOI: 10.1590/1982-0224-20170065

- Moore, A., Russel, I. C., & Potter, E. C. E. 1990. The effects of intraperitoneally implanted dummy acoustic transmitters on the behaviour and physiology of juvenile Atlantic salmon, *Salmo salar* L. *Journal of Fish Biology*, 37(5), 713-721. DOI: 10.1111/j.1095-8649.1990.tb02535.x
- Økland, F., Hay, C. J., Næsje, T. F., Thorstad, E. B., & Nickandor, N. 2001. Movements and habitat utilisation of radio tagged carp (*Cyprinus carpio*) in a reservoir in the Fish River, Namibia. NINA-NIKU Project Report, 13: 1-28.
- Ouedraogo, C., Canonne, M., D'Cotta, H., & Baroiller, J.-F. 2014. Minimal body size for tagging fish with electronic microchips as studied in the Nile tilapia. *North American Journal of Aquaculture*, 76(3): 275-280. DOI: 10.1080/15222055.2014.911228
- Schulz, U. H. 2003. Effects of surgically implanted dummy transmitters on the South American catfish Jundiá (*Rhamdia quelen*). *Brazilian Journal of Biology*, 63(2), 345-348. DOI: 10.1590/S1519-69842003000200019
- Soares, B. E., Barros, T. F., Hashiguti, D. T., Pereira, D. C., Ferreira, K. C. F., & Caramaschi, E. P. 2020. Traditional approaches to estimate length at first maturity (L50) retrieve better results than alternative ones in a Neotropical heptapterid. *Journal of Fish Biology*, 97(5), 1393-1400. DOI: 10.1111/jfb.14505
- Thorstad, E. B., Rikardsen, A. H., Alp, A., & Økland, F. 2013. The use of electronic tags in fish research – an overview of fish telemetry methods. *Turkish Journal of Fisheries and Aquatic Sciences*, 13(5), 881-896. DOI: 10.4194/1303-2712-v13_5_13
- Turek, K. C., Pegg, M. A., & Pope, K. L. 2014. Short-term evaluation of visible implant alpha tags in juveniles of three fish species under laboratory conditions. *Journal of Fish Biology*, 84(4): 971-981. DOI: 10.1111/jfb.12338
- Walsh, M. G., Bjorgo, K. A., & Isely, J. J. 2000. Effects of implantation method and temperature on mortality and loss of stimulated transmitters in hybrid striped bass. *Transactions of the American Fisheries Society*, 129(2), 539-544. DOI:10.1577/1548-8659(2000)129<0539:EOIMAT>2.0.CO;2
- Wilson, C. D., Arnott, G., Reid, N., Roberts, D. 2011. The pitfall with PIT tags: marking freshwater bivalves for translocation induces short-term behavioural costs. *Animal Behaviour*, 81(1), 341-346. DOI: 10.1016/j.anbehav.2010.10.003

Submitted: 04 January 2025

Accepted: 02 July 2025

Published: 19 September 2025

Associate Editor: Felipe Polivanov Ottoni