

Seasonal patterns in Gubernetes yetapa behavior

**SEASONAL SHIFTS IN BEHAVIOR AND SOCIAL
INTERACTIONS OF STREAMER-TAILED TYRANT (*GUBERNETES
YETAPA*) IN SOUTHERN BRAZIL**

Thuani Luísa Saldanha Wagener^{1*} , *Luis Eduardo Silva dos Santos*¹ , *Ramiro Silva
Almeida*² , *Nêmera Pauletti Prestes*³ , *Marilise Mendonça Krügel*⁴  & *Jaime Martinez*⁵ 

¹ Universidade Federal de Santa Maria, Programa de Pós-graduação em Biodiversidade Animal, Avenida Roraima, 1000, Camobi, CEP 97105-900, Santa Maria, RS, Brazil. huaniwagener@gmail.com

² Universidade Federal de Santa Maria, campus Palmeira das Missões. Avenida Independência, 3751 - Vista Alegre, CEP 98300-000, Palmeira das Missões, RS, Brazil. ³ Associação Amigos do Meio Ambiente, Departamento de Vida Silvestre, CEP 99500-000, Carazinho, RS, Brazil.

⁴ Universidade Federal de Santa Maria, Departamento de Engenharia Sanitária e Ambiental, Avenida Roraima, 1000, Camobi, CEP 97105-900, Santa Maria, RS, Brazil.

⁵ Universidade de Passo Fundo, Programa de Pós-graduação em Ciências Ambientais, BR 285 Km 292.7, São José, CEP 99052-900, Passo Fundo, RS, Brazil.

E-mails: thuaniwagener@gmail.com (*corresponding author); luis-santos.1@acad.ufsm; ramiroalmeida1@hotmail.com; prestes@upf.br; c; martinez@upf.br

SUPPLEMENTARY MATERIAL




Figure S1. Distribution of observation effort (%) throughout the day for two wetland populations (WL1 and WL2) of *Gubernates yetapa* in Palmeira das Missões, Rio Grande do Sul, Brazil. Bars represent the proportion of total sampling effort conducted during each hour, totaling 264 hours of field effort.




Figure S2. Frequencies of behavioral categories and their respective subcategories exhibited by *Gubernates yetapa* during 264 hours of field observations conducted in Palmeira das Missões, Rio Grande do Sul state. The breeding season was defined as October to February (dark blue bars), and the non-breeding season as March to September (grey bars).